

# Smallholder Farming Households Nutrition Under Extreme Heat: Vulnerabilities and Adaptation \*

Maxime Roche<sup>†</sup>

Version 18 November, 2025

[Latest version here](#)

## Abstract

Climate change threatens food security in Sub-Saharan Africa, but the magnitude of its nutritional impacts and the household-level behaviours that mediate them remain understudied. I examine how extreme heat during the crop growing season affects the commercialization and consumption decisions of Nigerian smallholder farming households, and how these responses shape post-harvest nutrition. Using five waves of panel data (2010–2024) linked to high-resolution weather, I exploit within-household variation in growing-season temperatures. Results show that hotter seasons depress harvests and reduce diet quality while leaving total caloric intake unchanged. I estimate that a  $+1^{\circ}\text{C}$  warming would result in 1.62 and 0.63 million additional households with inadequate protein and iron intake, respectively. Less educated households and those with young children, who require protein and iron for growth, are more adversely affected. I demonstrate that households reduce crop commercialization to safeguard calories from own-produced staples and cut their purchases of nutritious and cash-intensive foods. This reflects a risk-minimizing behaviour under incomplete food and labor markets and binding caloric constraints. In line with this, I find no evidence of adjustments in off-farm labor. Findings suggest that on-farm adaptation alone cannot maintain diet quality under climate change, underscoring the need for integrated policies that extend beyond calorie sufficiency and pair market participation and labor market development strategies with climate-risk mitigation.

**Keywords:** agriculture; climate change; extreme heat; Nigeria; nutrition.

**JEL Codes:** I15, O13, Q11, Q12, Q54.

---

\*I thank Laure de Preux, Olivier Ecker, Ariel Ortiz-Bobea, Franco Sassi, and participants to the Association of Environmental and Resource Economists 2025 and the European Association of Agricultural Economists 2025 conferences, IPWSD 2025 workshop at Columbia University, 10th DENEb workshop at the Potsdam Institute for Climate Impact Research (PIK), and seminar participants at Imperial College London for their helpful comments. This work was supported by funds from the UK Economic and Social Research Council through the London Interdisciplinary Social Science Doctoral Training Programme [grant number ES/P000703/1].

<sup>†</sup>Department of Economics & Public Policy, Imperial Business School, Imperial College London.  
[m.roche21@imperial.ac.uk](mailto:m.roche21@imperial.ac.uk)

# 1 Introduction

Undernutrition undermines human capital formation (Glewwe et al., 2001; Maluccio et al., 2009), deteriorates long-run health (Hoynes et al., 2016), and reduces productivity (Case and Paxson, 2008). The associated aggregate economic burden resulting from lost productivity and direct healthcare costs is substantial, at roughly 2-3% of global GDP each year (FAO, 2013). Despite notable progress over recent decades, the burden remains pronounced in Sub-Saharan Africa: undernourishment affects more than 22% of the population (FAO, 2022), and micronutrient deficiencies are widespread (FAO et al., 2021; Passarelli et al., 2024). At the same time, the region is warming faster than the global average (WMO, 2024). The increased frequency and intensity of extreme heat events could slow or even reverse progress against malnutrition (Blom et al., 2022). This is a salient issue for Sub-Saharan Africa, where the majority of households are smallholder farmers, for whom a hot growing season poses a direct threat to their primary sources of food and income.

While the link between extreme heat and reduced agricultural productivity is well-established (Hultgren et al., 2025; Schlenker et al., 2006; Schlenker and Roberts, 2009), much less is known about the pathways through which these productivity shocks translate into household nutrition. The magnitude of this impact is also uncertain, shaped by households' capacity to adapt. Faced with a harvest shortfall, a farm household's food consumption and crop commercialization decisions are intertwined under incomplete food and labor markets, common in such developing settings (Dillon and Barrett, 2017; Dillon et al., 2019). Yet, much of the literature has focused on coarse nutritional outcomes, such as self-reported food security or dietary diversity indices, which capture the extensive margin of consumption, or total energy intake (Dillon et al., 2015; Randell et al., 2022; Dasgupta and Robinson, 2022; Villacis et al., 2022; Kroeger, 2023). These metrics can mask important changes in dietary quality and obscure the commercialization–consumption nexus that mediates nutrition.

This paper examines the impact of extreme heat on food consumption and nutrition among smallholder farm households. I focus on Nigeria, the most populous African nation, with over 70% of households engaged in farming, high poverty rates, and widespread food insecurity (National Bureau of Statistics, 2024; World Bank, 2025). I study adjustments to consumption

and commercialization patterns in response to temperature-induced crop losses and assess their implications for undernutrition, diet quality, and nutrient intake.

I use five waves of nationally representative panel data from the Nigeria Living Standards Measurement Study - Integrated Surveys on Agriculture (LSMS-ISA), spanning 2010–2024. The survey provides microdata on household-level agricultural production, livestock management, and comprehensive food consumption diaries. I convert the latter into calories and the intake of crucial nutrients to dietary health, including protein, iron, zinc, and vitamin A, and create nutrient adequacy indicators based on international requirement standards and household composition. I then match these data with high-resolution gridded weather data. I model the relationship between weather and agricultural productivity as a non-linear function of cumulative exposure to heat and precipitation during the growing season. My hypothesis is that high growing-season temperatures increase crop losses, and that this decline in agricultural output is subsequently transmitted to reduced household welfare after harvest. The empirical strategy exploits quasi-random within-household variation in growing-season temperatures to identify the impact of extreme heat on post-harvest nutrition.

I find that hotter growing seasons significantly reduce harvests. In the short term, however, this productivity loss does not translate into a reduction in total energy intake. Instead, the shock's impact is on diet quality. Extreme heat leads to a decrease in household dietary diversity and an increase in the proportion of households with inadequate protein and iron intake. The magnitude is economically significant: a uniform  $+1^{\circ}\text{C}$  warming is associated with an additional 1.62 million and 0.63 million Nigerian households (4.04% and 1.56% of farm households) with inadequate protein and iron intake, respectively.

Tracing the mechanisms, I demonstrate that households respond to the harvest shortfall by adjusting their market behavior in a manner consistent with risk minimization under incomplete food and labor markets and binding caloric constraints. First, households sell a smaller share of their harvest. This adaptation allows them to maintain consumption of their own-produced foods, mainly cereals, roots, and tubers, thereby securing their caloric needs. Second, this decision, combined with the initial harvest loss, generates an income shock, reflected by a drop in total expenditure. Households cope by cutting spending on both food purchases and

non-food items. The cuts to food purchases disproportionately affect cash-intensive, nutritious foods such as vegetables, pulses, and animal meat, which are essential components of a healthy diet and significant sources of protein and iron. Thus, this *ex-post* adaptation strategy to ensure energy sufficiency directly degrades diet quality. Adverse nutritional effects worsen as time since harvest elapses and own-produced food stocks are depleted.

These adverse effects are not borne equally. The nutritional impacts are significantly larger for households with less-educated heads and those with young children, a concerning finding that aligns with evidence showing extreme heat increases the prevalence of child stunting (Blom et al., 2022). While closeness to markets and income diversity are associated with improved nutrition, they do not moderate the adverse effects of high growing-season temperatures. Livestock management, itself adversely affected by heat, does not buffer nutritional and income losses. Findings suggest that post-harvest market-level prices are largely unresponsive to growing season temperatures, at least in the short run. Consistent with incomplete labor markets, I find no statistically significant effects of extreme heat on post-harvest off-farm labor.

This paper makes three main contributions. First, it adds to the growing literature on heat's impact on nutrition in low-income settings with agricultural-dependent livelihoods (Dillon et al., 2015; Malacarne and Paul, 2022; Amare et al., 2021; Wegenast et al., 2025), by providing new causal evidence on the translation of heat shocks into malnutrition, moving beyond coarse, experience-based food security or energy-based metrics to focus on the intensive margin of nutrient adequacy. It shows that focusing on energy sufficiency can mask significant deteriorations in diet quality. While nutrient inadequacies - most notably protein and iron - are pervasive among Sub-Saharan African farm households, they have received limited attention in past economic analyses of food consumption, with the notable exception of McCullough et al. (2024).

Second, by separating consumption into own-produced and purchased sources, which is absent from the climate-nutrition literature, I provide evidence on how heat-induced harvest losses trigger defensive reallocation, consistent with caloric risk minimization under incomplete food and labor markets. Households retain a larger share of staple production for home use and cut market purchases of nutrient-dense foods, preserving calories at the expense of diet quality

in a pattern aligned with Bennett's law and the literature on farm households' consumption response to macroeconomic shocks (Headey et al., 2014; Ecker and Hatzenbuehler, 2022).

Third, I reveal an *ex-post* farm household adaptation response not previously documented. Following a heat shock in the growing season, households reduce commercialization and the share of harvest sold. The positive relationship between commercialization and smallholder farmers' nutrition is well-documented (Hazrana and Mishra, 2025; Chegere and Kauky, 2022; Ogutu et al., 2020). Increasing market participation through interventions that lower transaction costs, improve smallholders' access to markets and information, and strengthen producer organizations and rural infrastructure is essential to escaping semi-subsistence poverty traps (Omiti et al., 2009; Barrett, 2008). My findings suggest that climate change may represent an additional constraint to such participation. This complements a rich literature focusing on *ex-ante* productive responses to extreme weather shocks (Aragón et al., 2021; Mayorga et al., 2025; Costinot et al., 2016), the reallocation of economic activities (Rosenzweig and Stark, 1989; Kochar, 1999), and consumption smoothing techniques, such as asset sales and access to credit (Rosenzweig and Wolpin, 1993; Di Falco et al., 2011).

These findings highlight the limitations of on-farm strategies in mitigating nutritional risks under climate stress. Households' defensive adaptation preserves caloric intake but degrades nutrient adequacy, underscoring the need for policy approaches to extend beyond energy sufficiency. To protect nutrition, interventions must integrate climate-risk mitigation with strategies that expand income diversification and foster market participation.

This paper is divided as follows. [Section 2](#) develops a conceptual framework. [Section 3](#) details the Nigerian context and the data sources, including the construction of the main variables used in this study and summary statistics. [Section 4](#) describes the empirical strategy. [Section 5](#) presents the main results, investigates household adaptation responses, explores heterogeneity and potential moderators, and discusses the effect of a uniform warming scenario. Finally, [Section 6](#) concludes.

## 2 Conceptual framework

This section develops a conceptual framework to examine how smallholder farmers adjust commercialization decisions and diet composition in response to extreme heat. I depart from standard agricultural producer-consumer household models in the development literature (Benjamin, 1992; Janvry et al., 1991; Pitt and Rosenzweig, 1985), where households make simultaneous, non-separable decisions regarding consumption and commercialization.

Without loss of generality, I assume a one-person farm household framework with a single output,  $s$ , harvested after the growing season. I call this output “staples” and it can refer to any cereals, roots, or tubers commonly cultivated by Nigerian smallholder farmers.<sup>1</sup> Harvest for  $s$  depends on  $A$  (with  $s'(A) > 0$ ), a productivity shifter that captures the idea that farmers using identical inputs may face different levels of output due to farming skills, soil quality, or exposure to weather shocks.<sup>2</sup> Consistent with the agricultural yields and temperature literature (Hultgren et al., 2025), extreme heat ( $H$ ) has a detrimental effect on productivity, such that  $A'(H) < 0$ . Household’s utility is  $U(c, n)$ , where  $n$  is the consumption of non-staple “nutritious” foods (e.g., vegetables, fruits, animal-sourced foods), while  $c$  represents other market consumption. Staple consumption does not enter the utility function as they are a cheap way to meet basic calorie requirements, denoted  $\underline{K}$ , with  $k_s$  the calorie content of one unit of  $s$  and  $k_n$  the calorie content of one unit of  $n$ . The household obtains income by selling a share  $q \in [0, 1]$  of  $s$  or supplying off-farm working hours  $l \in [0, \bar{L}]$  at fixed wage  $w > 0$ .<sup>3</sup> I assume that the utility function is increasing and strictly concave and  $\frac{p_s}{k_s} < \frac{p_n}{k_n}$ .<sup>4</sup>

After each harvest season, the household maximizes utility by choosing off-farm labor  $l$ , the commercialization share  $q \in [0, 1]$ , selling  $qs$  and retaining  $(1 - q)s$  for home consumption, and the quantity of nutritious foods  $n$  and staple foods  $s_b$  purchased. In this simplified setting, the

---

<sup>1</sup>The National Bureau of Statistics (2024) shows that the most commonly cultivated crops among farming households (as a share of farming households growing crops) are maize (44.9%), sorghum (28.5%), millet (17.8%), and rice (16.2%) (cereals); cassava (44.9%) and yam (18.7%) (roots and tubers). Other crops, such as pulses or nuts, are less cultivated, e.g., beans (16.5%), groundnut (11.5%).

<sup>2</sup>I assume that capital is fixed, e.g., access to irrigation, which is low in Nigeria.

<sup>3</sup> $\bar{L}$  represents a cap on the number of hours the household can work off-farm in the post-harvest period. Because I define utility only over  $n$  and  $c$ , the model would otherwise push off-farm hours to infinity at any positive wage.  $\bar{L}$  is the maximum feasible hours given day length and other household and farm maintenance tasks.

<sup>4</sup>Globally, including in Sub-Saharan Africa, the price per kilocalorie is significantly lower for staples than nutritious foods (Masters et al., 2018; Bai et al., 2021).

farm household's problem is:

$$\begin{aligned} \max_{q,l} \quad & U(c, n) \\ \text{s.t.} \quad & p_s s_b + p_n n + c \leq p_s q s + w l \\ & k_s [(1 - q) s + s_b] + k_n n \geq \underline{K} \end{aligned}$$

If output and labor markets exist and are well functioning, consumption, commercialization, and labor decisions can be studied separately (Benjamin, 1992). In this case,  $p_s^{buy} = p_s^{sell} = p_s$  (i.e., no wedge); thus, commercialization is purely an accounting choice, and the household is indifferent between retaining their own-produced staples for consumption and selling them and then purchasing staples. A temperature-induced harvest shortfall represents an income shock, which can be offset by supplying more off-farm labor  $l \in [0, \bar{L}]$ . There is no sharp prediction on the effect on  $n$ , but intake should be maintained with ample earning capacity, i.e., high  $w$  or high  $\bar{L}$  compared to the initial off-farm labor supply endowment.

This prediction changes in the case of incomplete markets. Under incomplete food markets but complete labor markets, selling and buying staples destroy value as  $p_s^{buy} = p_s^{sell} + v_s$ . To meet calorie requirements cheaply after a harvest shortfall, the household avoids paying the wedge  $v_s > 0$  by retaining more of the smaller harvest. Cash needs for  $n$  are partially met by working more off-farm, and the overall impact on  $n$  depends on the earning capacity. Under incomplete labor markets (e.g., constrained by local demand) but complete food markets, there is no wedge on staple prices, and calories can be purchased without extra loss if there is cash. However, liquidity is tight because  $l$  cannot expand (or is limited). Therefore, the household sells a higher share of the harvest to raise cash when needed, and  $n$  declines, particularly when the calorie constraint is binding.

Finally, in the case with both incomplete food and labor markets, the cheapest way to secure calories is to retain  $s$ , thus  $\frac{dq}{dA} < 0$ , and the ability to earn cash is limited, thus  $\frac{dl}{dA} \approx 0$ . This result suggests that, in contexts with imperfect food and labor markets, extreme heat induces households to reduce commercialization and the consumption of nutritious foods. Because selling and repurchasing staples is costly in thin and incomplete markets, and post-harvest work

opportunities are rationed, households retain a larger share of the harvest to secure calories and scale back cash-intensive, nutritious food purchases. Households also face a risk of calorie shortfall if  $\underline{K}$  is high relative to  $s$ . The prediction is especially relevant where staple buy–sell wedges are large and local labor demand is limited, conditions that characterize many small-holder environments in Nigeria and similar developing settings (Dillon and Barrett, 2017; Dillon et al., 2019).

This framework also highlights forces that could confound a negative relation between extreme heat and commercialization. First, if extreme temperatures depress aggregate supply and raise output prices, higher prices would encourage sales, biasing the heat–commercialization coefficient toward zero or a positive value. I examine the role of prices in Section 5.6. Second, correlated productivity shocks can raise harvests and sales in periods that are also classified as hot, attenuating a negative effect. This could be the case with precipitations, thus I flexibly control for precipitation to net out correlated agronomic shocks.

Guided by this framework, the empirical analysis estimates the effect of extreme heat on home-grown consumption, commercialization, purchases, and subsequent dietary quality as measured both through dietary diversity and nutrient adequacy. Several limitations are worth noting. I focus on *ex-post* household adaptation, i.e., after the weather shock occurred (Carleton et al., 2024). At the time of commercialization and food consumption decisions, I treat harvest as predetermined, abstracting from *ex-ante* adaptation, such as production-stage adjustments. Post-harvest off-farm labor  $l$  does not affect production and thus enters only as an income-generation margin. The literature documents additional within-growing season responses, such as changes in input use (Mayorga et al., 2025; Aragón et al., 2021) and crop mix (Costinot et al., 2016), as well as consumption-smoothing strategies, including off-farm work and migration (Rosenzweig and Stark, 1989; Kochar, 1999). I examine these additional potential adaptation responses in Section 5.6.

## 3 Data

### 3.1 The Nigerian Context

Nigeria, the most populous nation in Sub-Saharan Africa with approximately 233 million people, serves as the setting for this analysis. The country faces considerable socioeconomic challenges, with 56% of its population living below the poverty line and over 30 million experiencing acute food insecurity (World Bank, 2025; World Food Programme and Food and Agriculture Organization of the United Nations, 2025). Agriculture is central to Nigeria's economy, involving over 70% of households (over 42 million). Major cultivated crops include maize, cassava, sorghum, yams, and millet (National Bureau of Statistics, 2024). Poverty is widespread among farm workers (World Bank, 2014). Agricultural productivity is hampered by several constraints, including limited market access, land degradation, insufficient infrastructure, and minimal irrigation (World Bank, 2025). Climate change intensifies these challenges and is projected to decrease agricultural productivity and worsen food insecurity, making Nigeria increasingly vulnerable (World Bank, 2021). Understanding farm household responses to rising temperatures is thus crucial and a key aspect of designing effective climate adaptation policies.

### 3.2 LSMS-ISA data

This study utilizes panel data from the Nigeria General Household Survey (GHS), part of the Living Standards Measurement Study-Integrated Surveys on Agriculture (LSMS-ISA). Specifically, it uses five waves collected in 2010/11 (wave 1), 2013/14 (wave 2), 2015/16 (wave 3), 2018/19 (wave 4), and 2023/24 (wave 5). The LSMS-ISA is a nationally representative and georeferenced household survey.<sup>5</sup> It captures detailed information on household demographics, socioeconomic characteristics, plot- and crop-level agricultural activities, livestock management, food prices, and food consumption. Demographic information is captured post-planting,

---

<sup>5</sup>A partial panel refresh was carried out in wave 4. Also, the representativeness of the GHS was impacted during wave 4 due to security issues, which prevented data collection in some locations. As a result, the wave 4 data is only representative of the areas that were safely accessible. The GHS does not track "split-off" households. I test the robustness of my results to the exclusion of the two latter waves 4 and 5 in Section 5.5.

while the other variables used in this analysis are derived from the post-harvest dataset collected between January and April.

A harmonised farm-to-fork micro panel dataset was constructed. Household-level agricultural output data were processed following the work of [Bentze and Wollburg \(2024\)](#). Only households engaged in crop cultivation, with available geolocation,<sup>6</sup> and observed for at least three waves were retained. Furthermore, households must have at least two waves with strictly positive agricultural output, plot area, and food consumption. This selection process yields a final sample of 9,006 observations from 2,832 unique households, distributed across 392 enumeration areas (clusters), representing 92.5% of the panel farm household-year observations from the five survey waves of the Nigeria LSMS-ISA ([Table C1](#)). [Figure 1](#) displays the geographical distribution of these clusters, illustrating the survey's comprehensive coverage across Nigeria.

Household food consumption data were collected through a 7-day recall, which detailed the food items consumed and their sources (own production, purchase, gift/transfer). It is important to note that these data reflect household-level food availability rather than individual intake, as they do not account for waste or intra-household distribution. Household food consumption data were processed following the work of [McCullough et al. \(2024\)](#). To assess nutritional status, food consumption quantities were harmonised to kilograms based on reported standard units or converted from non-standard units. Nutritional content was assigned using African food composition tables, adjusted for edible portions and potential cooking losses. Assumed fortification rates for relevant staples were incorporated, as described in [Appendix A](#). Household-level nutrient requirements - i.e., dietary energy, protein, iron, zinc, and vitamin A - were calculated based on the age and sex composition of the household, using estimated average requirements (EAR) from established sources and adult equivalence scales.<sup>7</sup> Adequacy percentages were constructed by dividing each household's total energy and nutrient availability by its EAR. Based on these percentages, dummy variables were created to indicate whether household-level caloric or nutri-

---

<sup>6</sup>Enumeration area coordinates are not publicly disclosed for wave 5. For this wave, I replaced missing coordinates with the enumeration area coordinates from the previous wave for the same enumeration area.

<sup>7</sup>For example, children and elderly members are counted as less than one full adult equivalent. See [Appendix A](#) for more details.




Figure 1. **Map of sample cluster locations.** Map of Nigeria, showing Northern (darker gray) and Southern (lighter gray) zones and regional borders. Cluster locations are represented by red dots. Based on 2,832 unique panel farm HH, within 392 enumeration areas (clusters).

ent consumption falls below 100, 80, or 60 percent of the recommended levels. The thresholds below 100% adequacy serve as a proxy to capture distributional impacts. Hereafter, I will refer to the 80% adequacy threshold as “suboptimal” intake and the 60% threshold as “insufficient” intake.

Finally, household dietary diversity is measured using the Household Dietary Diversity Score (HDDS), which reflects a household’s economic access to varied food and its food security. The standard HDDS includes 12 food groups (FAO, 2011).<sup>8</sup> Following Nguyen and Qaim (2025), the oils/fats, sugar/honey, and miscellaneous food groups are excluded, as these are often considered less nutritious and may skew the HDDS as a measure of dietary quality

<sup>8</sup>The 12 food groups are: cereals; roots & tubers; vegetables; fruits; meat, poultry, and offal; eggs; fish and seafood; pulses, legumes, and nuts; milk and milk products; oil/fats; sugar/honey; miscellaneous (mostly spices, condiments, and beverages) (FAO, 2011).

	Mean	Median	SD	Min.	Max.
Head age (years)	52.87	52.00	14.62	16.00	100.00
Head formal education (1/0)	0.40	0.00	0.49	0.00	1.00
Head female (1/0)	0.14	0.00	0.35	0.00	1.00
Child < 5 yo (1/0)	0.47	0.00	0.50	0.00	1.00
Child < 15 yo (1/0)	0.79	1.00	0.41	0.00	1.00
HH size	6.06	6.00	3.25	1.00	31.00
Total yield (kg/ha)	8,224.58	3,063.20	12,837.06	0.00	56,497.18
Total harvest (kg)	3,181.84	1,500.00	6,255.99	0.00	161,000.00
Total area (ha)	1.00	0.51	1.64	0.00	42.57
Share of harvest sold	0.20	0.00	0.30	0.00	1.00
HH livestock under management (1/0)	0.69	1.00	0.46	0.00	1.00
HH annual total exp per capita (USD 2020)	671.77	490.24	956.84	43.97	38,805.34
HH dietary diversity score (HDDS)	5.58	6.00	1.61	1.00	9.00
HH own-production share of energy intake	0.46	0.49	0.28	0.00	1.00
Observations	9,006				

Table 1. **Household descriptive statistics.** Agricultural output values are winsorized at the 99th percentile. Annualized expenditure in USD 2020 (originally, food: last 7 days; non-food: last month or last 12 months). Expenditure information is missing for wave 5. Household dietary diversity score (HDDS) is defined as nine groups based on (Nguyen and Qaim, 2025). HH: household.

(Verger et al., 2019).

Table 1 provides descriptive statistics on household characteristics and agricultural output. On average, households have a total plot area of 1.0 hectare (ha). The majority of households do not participate in the commercialization of their harvest. Over two-thirds of the sample manage livestock. Average daily per capita expenditure is USD 1.84 (2020 USD), reflecting the high prevalence of extreme poverty among Nigerian farm households. The median household reports having consumed six out of the nine food groups that comprise the HDDS in the last seven days, with cereals and vegetables representing the two groups with the highest likelihood of consumption (over 97%) (Figure B1).

Figure 2 displays the average share of sample households with adequate dietary intake for energy, protein, and three micronutrients, including iron, zinc, and vitamin A. Protein is crucial for basic growth and tissue repair. The selected micronutrients are critical for vital physiological functions: iron for oxygen transport (preventing anemia), vitamin A for vision and immunity, and zinc for immune function and growth (Passarelli et al., 2024). Deficiencies in these areas lead to severe outcomes like stunted growth, weakened immunity, and cognitive impairment,




Figure 2. **Average energy and nutrient adequacy levels.** HH: household.

all of which hinder productivity and development (Black et al., 2013). The mean prevalence of household energy adequacy is 42%. Among nutrients, adequacy is highest for vitamin A (87%) and lowest for zinc (54%). As expected, the share of sample households with adequate dietary intake is higher for lower adequacy levels.

### 3.3 Weather data

Household locations (cluster centroids) were matched with high-resolution gridded weather data. Daily temperature data were obtained from the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 reanalysis dataset, which provides data at a  $0.1^\circ \times 0.1^\circ$  resolution (approximately  $10\text{km} \times 10\text{km}$ ) (Hersbach et al., 2020). Precipitation data were derived from the University of California, Santa Barbara Climate Hazards Center Infrared Precipitation with Station (CHIRPS) dataset (Funk et al., 2015). Originally at  $0.05^\circ \times 0.05^\circ$  resolution, monthly precipitation data were reaggregated using means to match the  $0.1^\circ \times 0.1^\circ$  ERA5 grid. The weather data for a specific grid square was assigned to an LSMS-ISA cluster if the cluster lay within that grid square.

The growing season is defined based on the Famine Early Warning Systems Network (FEWS-




Figure 3. **Temperature distribution of growing season days.** Based on the sample farm panel households, i.e., 9,006 observations, 2,832 HH, within 392 enumeration areas (clusters), over 2010-2023.

NET) agricultural calendar for Nigeria (Figure B5).<sup>9</sup> It refers to the period during which crops are planted and grown. It is defined as March to August for the Southern region (including the South East, South South, and South West administrative regions) and May to September for the Northern region (including the North Central, North East, and North West administrative regions). This definition aligns with the distribution of the percentage of planted plots per month derived from the 39,030 plots cultivated by the sample farm households (Figure B6). Figure 3 shows the distribution of temperatures observed during the last completed growing season for the whole sample.<sup>10</sup>

<sup>9</sup>Source: [Famine Early Warning Systems Network](#), Season Calendar (Accessed 10 April 2025).

<sup>10</sup>Figure B7 presents a similar distribution by region.

## 4 Empirical strategy

The empirical analysis aims to identify how households respond to extreme heat and its impact on household nutrition. The primary channel through which weather affects farm households is agricultural production. Assuming farm-level production in Nigeria can be represented by a Cobb-Douglas production function,<sup>11</sup> where  $y_{i,t}$  denotes agricultural output (post-harvest) for household  $i$  following growing season  $t$ , weather conditions in the growing season affect output production through their effect on the productivity term  $A_{i,t}$  (Aragón et al., 2021). I approximate the reduced-form nutritional impact of weather conditions during the last growing season using the following log-linear regression model:

$$y_{i,t} = g(\beta, \mathbf{w}_{i,t}) + \lambda_m + \mu_t + \eta_i + \varepsilon_{i,t} \quad (1)$$

where the  $y$  is post-harvest agricultural output or a nutritional outcome and  $g(\beta, \mathbf{w}_{i,t})$  is a non-linear function of temperature and precipitation  $\mathbf{w}_{i,t}$  during the last growing season. The parameter of interest is  $\beta$ , which represents the vector of reduced-form estimates of the effect of weather shocks on farm household nutrition. This multi-way panel fixed effects regression approach is common in the climate economics literature (Dell et al., 2014). The identification strategy leverages the quasi-random variation in weather exposure within-households across growing seasons. I include a comprehensive set of fixed effects to isolate this exogenous variation. The terms  $\lambda_m$ ,  $\mu_t$ , and  $\eta_i$  are respectively a set of month-of-interview fixed effects, growing season fixed effects, and household fixed effects.  $\lambda_m$  control for seasonality in nutrition, such as the depletion of food stocks as the interview date gets further from the last harvest.  $\mu_t$  accounts for any country-wide changes in weather conditions or nutrition that may be spuriously correlated over time, as well as macro-level trends and systemic measurement error across survey waves.  $\eta_i$  controls for unobserved time-invariant household characteristics and heterogeneity.

Based on the seminal work of Schlenker et al. (2006), I model the nonlinear relationship between temperature and productivity using a degree-days approach. This method allows for

---

<sup>11</sup> $y_{i,t} = A_{i,t}K_{i,t}^{\alpha_1}L_{i,t}^{\alpha_2}X_{i,t}^{\alpha_3}$ , where  $A_{i,t}$  represents the productivity term,  $K_{i,t}$  capital use (e.g., land area),  $L_{i,t}$  labor use (e.g., family and hired labor), and  $X_{i,t}$  input use (e.g., fertilizer).

temperature exposure effects to be cumulative, with temperature exposure being beneficial up to a certain point and harmful thereafter. Specifically, I construct two measures for each household  $i$  in growing season  $t$ : growing degree days ( $GDD_{i,t}$ ) and harmful degree days ( $HDD_{i,t}$ ).  $GDD_{i,t}$  capture the cumulative exposure to temperatures between  $8^\circ\text{C}$  and a threshold  $\tau$ ,<sup>12</sup> while  $HDD_{i,t}$  captures exposure to extreme heat, i.e., temperatures above  $\tau$ . Formally:

$$\begin{aligned} GDD_{i,t} &= \sum_{d=1}^n (\min\{h_{i,d}, \tau\} - 8) \mathbf{1}\{h_{i,d} \geq 8\}, \\ HDD_{i,t} &= \sum_{d=1}^n (h_{i,d} - \tau) \mathbf{1}\{h_{i,d} > \tau\}. \end{aligned} \tag{2}$$

Here,  $h_d$  is the average temperature on day  $d$  and  $n$  is the total number of days in the growing season. A key empirical challenge is defining  $\tau$ , the temperature threshold above which harm occurs. This threshold is likely crop- and context-specific. Estimates from studies in other settings may not be transferable due to differences in crop mix, agricultural technology, and climate. Therefore, I adopt a data-driven approach by estimating a series of regressions of log agricultural output on  $GDD_{i,t}^\tau$  and  $HDD_{i,t}^\tau$ , iterating  $\tau$  across a range of plausible temperatures, similar to the procedure used in [Schlenker et al. \(2006\)](#). I then select the optimal threshold,  $\tau^*$ , that provides the best model fit (i.e., the highest  $R^2$ ).

I control for total growing season precipitation ( $P_{i,t}$ ) and its square to account for the cumulative non-linear effects of rainfall and avoid omitted-variable bias.<sup>13</sup> This leads to the following parametrization of the function  $g(\beta, \mathbf{w}_{i,t})$ :

$$g(\beta, \mathbf{w}_{i,t}) = \beta_1 GDD_{i,t}^{\tau^*} + \beta_2 HDD_{i,t}^{\tau^*} + \beta_3 P_{g,y} + \beta_4 P_{g,y}^2 \tag{3}$$

The dependent variable in [Equation 1](#) is transformed into logarithms for continuous variables. This is problematic for the food group-level analysis for a subset of households that reported zero consumption for specific food groups ([Figure B1](#)). To avoid the issue of an un-

---

<sup>12</sup> $8^\circ\text{C}$  is a common lower bound for growing degree days in the literature ([Arag3n et al., 2021](#); [Schlenker et al., 2006](#); [Schlenker and Roberts, 2009](#)).

<sup>13</sup>The main practical risk is multicollinearity, as temperatures and precipitations can move together at seasonal scales (e.g., during the growing season), which can inflate standard errors and make the separate coefficients less precise. However, this impacts precision, not consistency.

defined variable, I adopt the solution proposed by [Blakeslee and Fishman \(2018\)](#) and [Carpena \(2019\)](#). Specifically, zero-consumption observations are handled by setting the dependent variable to zero and simultaneously including a dummy variable to account for this data adjustment in the regression.<sup>14</sup>

Lastly, I account for potential spatial and temporal correlation in the error terms ( $\varepsilon_{i,t}$ ) by estimating spatial heteroskedasticity and autocorrelation Consistent (HAC) standard errors based on [Conley \(1999\)](#), using a distance parameter of 100 km and a temporal lag parameter of 5 years.<sup>15</sup>

## 5 Main results

### 5.1 Temperatures and agricultural production

This section presents my main empirical results on the effect of extreme heat on farmers' productivity, as measured by total harvest (in kilograms). First, [Figure 4](#) documents the outcomes from the iterative procedure consisting of nine regressions with different threshold  $\tau$  ranging from 26°C to 34°C and the comparison of model fit. This procedure yields an optimal threshold of  $\tau^* = 31^\circ\text{C}$ .<sup>16</sup> I use *GDD* and *HDD* calculated with this threshold in all subsequent regressions.

[Table C3](#) presents the results obtained from estimating [Equation 1](#) on total yield, harvest, and land area. Consistent with previous findings in other contexts ([Schlenker and Roberts, 2009](#); [Hultgren et al., 2025](#); [Burke and Emerick, 2016](#)), the estimates suggest that extreme heat has a negative effect on agricultural productivity and output. Each additional HDD in the last completed growing season results in a 1.54% decrease in total harvested quantities. To put this

---

<sup>14</sup>I provide a robustness check measuring household energy intake at the food group level in levels rather than logarithms in [Section 5.5](#).

<sup>15</sup>Given that my choice of distance and temporal lag parameters of the spatial HAC standard errors is arbitrary, I assess the sensitivity of my main findings to variations in these parameters in [Section 5.5](#).

<sup>16</sup>[Figure B8](#) presents the results for a similar iterative procedure using total yield (kg/ha) as outcome, yielding the same threshold  $\tau^* = 31^\circ\text{C}$ .




Figure 4. **Optimal GDD/HDD threshold using an iterative regression approach and effect on total harvest.** The left panel compares model fit (measured as  $R^2$ ) between nine regressions, one for each GDD/HDD threshold between 26°C to 34°C. The right panel compares the marginal impact of one HDD for each of the nine regressions. Total harvest is measured in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional growing season HDD. Conley spatial HAC standard errors (in parentheses).

figure in perspective, note that under a uniform +1°C warming scenario, [Section 5.7](#) projects that the average number of HDD during the growing season could increase by 12.95.

## 5.2 Temperatures, dietary diversity, and the intake of energy and nutrients

I estimate [Equation 1](#) to test how extreme heat impacts the intake of energy and nutrients. Despite the extreme heat’s impact on total harvested quantities, it does not have a statistically significant impact on average energy and nutrient intake.<sup>17</sup> However, I find that extreme heat during the last completed growing season decreases household dietary diversity, with statistical significance at the 10% level ([Table 2](#)), in line with [Dillon et al. \(2015\)](#)’s findings using cross-sectional data from wave 1 of the Nigeria LSMS-ISA panel. This effect is driven by a reduced likelihood of consuming nutritious foods, including fruits, meat, and eggs ([Figure B9](#)). These

<sup>17</sup>[Table C4](#) shows similar results considering the effect of extreme heat on the average intake of the macronutrient components of total energy, namely carbohydrates, protein, and fat.

	HDDS	Energy	Protein	Iron	Zinc	Vitamin A
GDD in growing season	-0.0005 (0.0003)	0.0001 (0.0002)	0.0001 (0.0002)	0.0002 (0.0002)	0.0003 (0.0003)	0.0003 (0.0002)
HDD in growing season	-0.0053* (0.0029)	-0.0002 (0.0015)	-0.0005 (0.0015)	0.0004 (0.0017)	0.0007 (0.0016)	0.0016 (0.0021)
Household FE	Yes	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.597	0.554	0.605	0.620	0.516	0.455
Observations	9,006	9,006	9,006	9,006	9,006	9,006

Table 2. **Temperatures, dietary diversity, and the intake of energy and nutrients.** The household dietary diversity score (HDDS) is defined as nine food groups based on [Nguyen and Qaim \(2025\)](#). Coefficients on HDDS can be interpreted as the absolute change in the HDDS for one additional HDD (sample mean: 5.58). Total energy, protein, iron, zinc, and vitamin A are measured in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Conley spatial HAC standard errors (in parentheses). FE: fixed effects.

food categories, which represent essential components of a diverse diet and rich sources of the nutrients under study, are also among the least frequently consumed by sample households ([Figure B1](#)).

Next, I test if extreme heat affects the number of households with adequate energy intake. [Figure 5](#) presents the estimates for the marginal effect of one additional HDD in the last growing season on the percentage of households with energy intake above 100%, 80%, and 60% adequacy. It shows that there is no statistically significant effect of extreme heat on the percentage of households that consume 100% or 80% of their energy requirements. In contrast, I find that an additional HDD in the last growing season increases the percentage of households with insufficient caloric intake - defined as not meeting at least 60% of their energy requirements - by 0.16 percentage points, however, this increase is not statistically significant at conventional levels ( $p = 0.112$ ). These results suggest that extreme heat could exacerbate undernutrition for households already experiencing it at extreme levels.

[Figure 5](#) shows the impact of extreme heat on household nutrient adequacy. Hot growing seasons negatively affect protein and iron adequacy among sample households. For protein, one additional HDD in the prior growing season increases the percentage of households below the 100% and 80% adequacy thresholds by 0.26 and 0.22 percentage points, respectively. For




Figure 5. **Effect of HDD on nutrient adequacy.** Each nutrient adequacy represents a dummy equal to one if a household meets the intake requirement at various thresholds (60%, 80%, and 100%). Coefficients have been divided by the respective dependent variable sample mean, and can be interpreted as the relative percentage change in the likelihood of meeting the requirement for one additional HDD. Conley spatial HAC standard errors are used to build the 95% confidence intervals.

iron, it increases the percentage of households below the 80% and 60% adequacy thresholds by 0.25 and 0.19 percentage points, respectively. The effect sizes are smaller for the 60% nutrient adequacy threshold. This can be explained in part by the smaller share of households near or below this threshold (Figure 2). Lastly, I find no evidence that extreme heat in the last growing season pushes households below nutrient adequacy levels for zinc and vitamin A intake, at any of the considered levels.

In Figure B10, I examine what is driving these nutrient adequacy results by estimating the effect of extreme heat on energy intake by food group. The post-harvest intake of staple foods, including cereals, roots and tubers, and oil/fats, is not affected by extreme heat in the growing season. These food groups represent the three most important sources of energy for the average Nigerian farm household, with 54.8% for cereals, 17.1% for root & tubers, and 12.3% for oil/fats (Figure B2). This participates in explaining the null effect of extreme heat on total energy intake in Table 2. On the other hand, an additional HDD in the last growing season reduces the post-harvest intake of more nutritious foods, which are important sources of

protein and iron, such as vegetables (-0.68%), meat (-0.33%), and pulses (-0.38%), as well as fruits (-0.30%).

### 5.3 Farm household response: commercialization and food purchase

As shown in [Figure 4](#) and [Table C3](#), farm households experience a negative shock to their harvest following extreme heat during the growing season. I am interested in examining their *ex-post* (i.e., post-harvest) adaptation responses and test the hypothesis laid out in [Section 2](#). First, in [Table C5](#), I examine changes in total energy intake, by source of consumption, i.e., purchased, own-produced, or gifted. Purchased foods represent approximately half of the total energy intake for the average household in the sample (49.5%), while own-produced foods represent 46.4%, and the rest is received as assistance or gift (4.1%) ([Table 1](#)). Extreme heat has no statistically significant effects on total energy intake from any consumption source, with a negative sign for the impact of one additional HDD on total energy purchased. Second, in [Figure 6](#), I examine changes in purchased and own-produced calorie intake separately for each food group. Extreme heat has no statistically significant effects on energy intake from own-produced food, except for fruits. On the other hand, an additional HDD in the last growing season reduces energy intake from purchased vegetables (-0.75%), meat (-0.30%), and pulses (-0.49%).<sup>18</sup> These results suggest an adjustment toward meeting energy sufficiency through own production, mainly composed of staple cereals, roots, and tubers, rather than purchasing more nutritious foods, thereby worsening dietary quality.

Two pathways can support an increased post-harvest reliance on own-produced food following a negative agricultural shock. First, households could reduce commercialization. Second, households could modify their own-produced food consumption smoothing behavior by prioritizing own-produced food intake immediately post-harvest, thereby risking the faster depletion of these resources.

[Table 3](#) reports the effect of extreme heat on commercialization and household expenditure. In line with the hypothesis in [Section 2](#) and the above-mentioned first potential pathway behind

---

<sup>18</sup>Purchased energy represents the majority of total energy intake from vegetables (84.2%), meat (88.0%), and pulses (66.5%) ([Figure B4](#)).




Figure 6. **Effect of HDD on energy intake, by food group and source of consumption.** Each food group is expressed as the logarithm of total energy intake, thus coefficients can be interpreted as the relative percentage change for one additional HDD. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors are used to build the 95% confidence intervals.

increased reliance on own-produced food post-harvest, I find that one additional HDD in the last growing season has a negative impact on both the extensive and intensive margins of commercialization. It decreases the likelihood of selling at least some harvested quantities by 0.40 percentage points (or relatively by 0.99%) and the share of total harvest sold by 0.18 percentage points (relatively by 0.90%).

In line with findings by others in the literature (Carpena, 2019; Dillon et al., 2015), Table 3 also shows that the drop in crop production is accompanied by lower household spending, likely aggravated by reduced commercialization as a share of total harvest. Indeed, all HDD coefficients are negative and statistically significant. Total expenditure, a proxy for income in the development economics literature (Carletto et al., 2021), decreases by 0.74% for each additional growing season HDD.<sup>19</sup> Extreme heat also decreases food purchases. The higher income

<sup>19</sup>Total expenditure includes a valuation of own-produced consumption, reducing classical measurement error and capturing resources actually available for nutrition.

	Total exp	Food exp	Non-food exp	Seller	Share sold
GDD in growing season	-0.0008*** (0.0003)	-0.0001 (0.0002)	-0.0006*** (0.0001)	-0.0004*** (0.0001)	-0.0002*** (0.0001)
HDD in growing season	-0.0074*** (0.0017)	-0.0034** (0.0016)	-0.0075*** (0.0014)	-0.0040*** (0.0011)	-0.0018*** (0.0006)
Household FE	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.661	0.591	0.749	0.471	0.470
Observations	8,146	8,146	8,146	5,970	5,970
Mean Y	.	.	.	0.405	0.199

Table 3. **Temperatures, expenditure, and commercialization.** Expenditure is expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Annualized expenditure in USD 2020 (originally, food: last 7 days; non-food: last month or last 12 months). Food expenditure represents purchased food. Total expenditure includes a valuation of own-produced consumption. Expenditure information is missing for wave 5. Seller is a dummy variable equal to one if a household is selling any harvested quantities. Information on sold harvested quantities is missing for 33.7% of households (either missing or set to missing because it is higher than harvested quantities or negative). I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors (in parentheses). exp: expenditure. FE: fixed effects.

elasticity of the demand for more nutritious foods, particularly vegetables and meat, compared to staples (Colen et al., 2018), explains the targeted cuts in purchases for these foods (Figure 6). This pattern is consistent with Bennett’s law operating in reverse: when resources contract, diets de-diversify away from purchased nutritious foods toward staples (Headey et al., 2014; Ecker and Hatzenbuehler, 2022).

Additionally, it is important to understand how extreme heat impacts non-food expenditures. Non-food spending is often a reliable indicator of a household’s economic well-being (Deaton, 1997). It also has consequences for a household’s food budget. This relationship could go in two directions. Households might cut back on non-food items to protect food purchases, or, alternatively, non-food purchases could end up limiting the funds available for food purchases. I show that one additional growing season HDD leads to a 0.75% decrease in non-food expenditure, supporting the former. As can be seen in Table C6, extreme heat affects most types of non-food spending, including housing, communication and transport, and clothing. The largest negative effects are observed for housing, where one additional HDD in the previous growing

season results in a 0.94% decrease in expenditure. Housing includes expenses on utilities (water, electricity, gas, fuels, and refuse collection), maintenance, domestic household services, and rent. Although the HDD coefficient is negative for education, it is not statistically significant at conventional levels.

Next, I examine the impact of the uncovered post-harvest household risk-minimizing strategy, which consists of prioritizing own-produced food intake post-harvest to ensure energy sufficiency, on consumption smoothing. For this, I estimate the heterogeneous effect of one additional HDD by survey months. Interviews for the Nigeria LSMS-ISA post-harvest household questionnaire are conducted between January and April, with the majority taking place in February and March.<sup>20</sup> Table 4 shows that the negative impact of extreme heat on dietary diversity, as well as protein and iron suboptimal adequacy (80%), intensifies with time since harvest.<sup>21</sup> This is likely driven by both the depletion of the income collected from the sale of harvest, as illustrated by a stronger reduction in food purchases in March-April compared to January-February, and the depletion of home-grown food reserves with a decline in own-produced food intake in March-April (Table C7). The latter is caused by a lesser energy intake from own-produced cereals, vegetables, and pulses (Table C8). This results in a more adverse impact on insufficient energy adequacy (60%), however, non-statistically significant at conventional levels (Figure B12).

#### 5.4 Heterogeneity and potential moderators

I explore heterogeneity in the effect of the last growing season temperatures on household nutrition to identify the most vulnerable populations. Table 5 presents heterogeneity by head sex and education, as well as the presence of children below five years old in the household. Results suggest that the effect of HDD on dietary diversity and nutrient adequacy is weaker among households with heads with formal education. On the other hand, the impact of extreme heat on malnutrition is qualitatively similar between households with and without a female head. Formal education could be correlated with a higher ability to participate in non-farm

---

<sup>20</sup>Only the post-harvest household questionnaire for wave 4 is collected in January (Table C2). Thus, I conduct a robustness check omitting wave 4 (Table C9), finding similar results.

<sup>21</sup>Figure B12 presents the results for this heterogeneity analysis by adequacy threshold for each nutrient.

	HDDS	Adq prot	Adq iron	Food exp	Non-food exp	Share sold
GDD	-0.0003 (0.0004)	0.0001 (0.0001)	0.0001 (0.0001)	0.0001 (0.0002)	-0.0006*** (0.0001)	-0.0002** (0.0001)
GDD × Mar-Apr	-0.0002 (0.0001)	-0.0002*** (0.0000)	-0.0002*** (0.0001)	-0.0003*** (0.0001)	-0.0000 (0.0001)	-0.0000 (0.0000)
HDD	-0.0017 (0.0030)	-0.0012 (0.0009)	-0.0015 (0.0009)	-0.0024 (0.0016)	-0.0074*** (0.0013)	-0.0017*** (0.0006)
HDD × Mar-Apr	-0.0047*** (0.0015)	-0.0011*** (0.0004)	-0.0011** (0.0004)	-0.0016** (0.0007)	0.0004 (0.0006)	0.0001 (0.0003)
Household FE	Yes	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.598	0.484	0.493	0.620	0.749	0.471
Observations	9,006	9,006	9,006	8,146	8,146	5,970
Mean Y	5.576	0.829	0.750	.	.	0.199

Table 4. **Temperatures, dietary diversity, nutrient adequacy, expenditure, and commercialization, by survey months.** Adequacy indicators are set to 80% adequacy levels. Coefficients on HDDS can be interpreted as the absolute change in the HDDS for one additional HDD. Coefficients on adequacy indicators can be interpreted as percentage-point change in the likelihood for one additional HDD. Expenditure is expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Annualized expenditure in USD 2020 (originally, food: last 7 days; non-food: last month or last 12 months). Food expenditure represents purchased food. Expenditure information is missing for wave 5. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Information on sold harvested quantities is missing for 33.7% of households (either missing or set to missing because it exceeds the harvested quantities or is negative). Conley spatial HAC standard errors (in parentheses). Adq: Adequacy. FE: fixed effects. HDDS: Household dietary diversity score.

employment. However, incomplete labor markets may limit the availability of such non-farm opportunities (Dillon et al., 2019). I investigate this further in Section 5.6. A complementary, compositional channel is that education is positively associated with household diet quality net of resources (Rashid et al., 2011). Mechanistically, schooling could attenuate the impact of shocks by enhancing nutritional knowledge, inventory management, or financial planning to smooth consumption when extreme heat tightens resource constraints. Lastly, the nutrition of farm households with young children, who require protein and iron for growth (Black et al., 2013), is the most adversely impacted. This finding is concerning, given that over 30% of children below five years old remain stunted in Nigeria (Akombi et al., 2017).

It is interesting to investigate potential moderators of the adverse effects of extreme heat on dietary diversity and nutrient adequacy to inform strategies for improving resilience to cli-

	HDDS	Adq prot	Adq iron	Food exp	Non-food exp	Share sold
HDD	-0.0050* (0.0028)	-0.0021*** (0.0008)	-0.0023*** (0.0008)	-0.0039** (0.0016)	-0.0067*** (0.0014)	-0.0018*** (0.0006)
HDD × Head formal edu	0.0038*** (0.0014)	0.0010*** (0.0003)	0.0011*** (0.0004)	0.0021*** (0.0006)	-0.0003 (0.0006)	0.0003 (0.0003)
HDD	-0.0053* (0.0028)	-0.0022*** (0.0008)	-0.0024*** (0.0009)	-0.0042*** (0.0016)	-0.0075*** (0.0014)	-0.0017*** (0.0006)
HDD × Head female	0.0041 (0.0027)	0.0004 (0.0007)	-0.0004 (0.0007)	0.0034** (0.0015)	0.0011 (0.0014)	0.0000 (0.0007)
HDD	-0.0025 (0.0028)	-0.0019** (0.0008)	-0.0019** (0.0009)	-0.0025 (0.0017)	-0.0080*** (0.0014)	-0.0014** (0.0006)
HDD × Age < 5 yo	-0.0044*** (0.0013)	-0.0005* (0.0003)	-0.0010** (0.0004)	-0.0024*** (0.0007)	0.0006 (0.0006)	-0.0006* (0.0003)
Mean Y	5.576	0.829	0.750	.	.	0.199

**Table 5. Effect of HDD on dietary diversity, nutrient adequacy, expenditure, and commercialization, by household characteristics.** This table displays the results from 16 regressions, under the main specification. Adequacy indicators are set to 80% adequacy levels. Coefficients on HDDS can be interpreted as the absolute change in the HDDS for one additional HDD. Coefficients on adequacy indicators can be interpreted as percentage-point change in the likelihood for one additional HDD. Expenditure is expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Annualized expenditure in USD 2020 (originally, food: last 7 days; non-food: last month or last 12 months). Food expenditure represents purchased food. Expenditure information is missing for wave 5. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Information on sold harvested quantities is missing for 33.7% of households (either missing or set to missing because it is higher than harvested quantities or negative). Conley spatial HAC standard errors (in parentheses). Adq: Adequacy. edu: education. HDDS: Household dietary diversity score.

mate change. Closeness to markets, livestock management, and income diversity have all been found to be correlated with improved nutrition in Sub-Saharan Africa (Babatunde and Qaim, 2010; Nguyen and Qaim, 2025; Headey et al., 2018). I examine these potential moderators by estimating heterogeneous responses to extreme heat. I interact HDD with the logarithm of the distance in kilometers to the nearest population center with more than 20,000 inhabitants. This represents a proxy for access to food markets in the literature, both for selling harvest and purchasing food (Nguyen and Qaim, 2025). I conduct similar interactions with three dummy variables: one equal to one if the household managed at least one livestock head in the last 12 months, one equal to one if the household ran at least one non-farm enterprise in last 12

months, and one equal to one if the household head was employed in a wage job in the last seven days before the post-harvest interview. Results suggest no moderating effects on nutritional indicators for livestock management and non-farm enterprise. A higher distance to the closest population center is negatively associated with the effect of an additional HDD on the likelihood of a household meeting suboptimal iron adequacy (80%), although this association is only statistically significant at the 10% level. A post-harvest wage job is positively associated with the effect of an additional HDD on the likelihood of a household meeting suboptimal iron adequacy and on the share of harvest sold, suggesting that off-farm employment may play a buffering role, reducing the need for households to pursue the risk-minimization strategy highlighted in [Section 5.3 \(Table C10\)](#).

I investigate the non-moderating role of livestock management further. Managing livestock could serve as both a nutritional buffer, as animal-sourced foods represent a source of quality protein among farm households, and an income buffer through the sale of livestock and their products ([Headey et al., 2018](#); [Rosenzweig and Wolpin, 1993](#)). First, I estimate the effect of extreme heat on the number of livestock heads managed by panel farm households involved in livestock over the last 12 months.<sup>22</sup> Given the year-round exposure of livestock and the lack of a well-defined ‘growing temperature range’ for animals, I adopt a binned temperature approach rather than HDD. This allows me to flexibly estimate the nonlinear effects of discrete temperature extremes on herd outcomes over the last 12 months.<sup>23</sup> Replacing a day with an average temperature  $< 31^{\circ}\text{C}$  by a day  $> 35^{\circ}\text{C}$  in the last 12 months reduces the number of large ruminant heads under management (bulls, cows, steers, heifers, calves, or ox) by 7.22%, with no statistically significant effects on the number of small ruminant heads under management

---

<sup>22</sup>I follow the same data processing criteria as for panel farm households, as described in [Section 3](#). Only households engaged in livestock management and observed for at least three waves were retained. Furthermore, households must have at least two waves with strictly positive livestock heads. This selection process yields a final sample of 5,627 observations from 1,949 unique households, accounting for 62.48% of the panel farm household-year observations across the five survey waves of the Nigeria LSMS-ISA.

<sup>23</sup>I depart from the degree-days specification used elsewhere in the paper and instead use temperature bins for the livestock analysis for two reasons. First, the outcome variable (i.e., number of livestock heads) is measured as an annual stock, not tied to a growing season. A 12-month exposure window, combined with livestock’s physiological response to extreme heat, is likely to exhibit nonlinear effects at extreme temperatures. Second, unlike crops, livestock lack a defined “growing” temperature range. Using temperature bins allows for easier interpretation of high-heat exposure episodes across the full year, rather than relying on cumulative heat intensity as in HDD, which is better suited to crop yield modeling.

(goats or sheep). Days with average temperature between 31°C and 35°C have no statistically significant effects (Figure 7).

Figure B11 shows that this result is driven by increased death or losses of large ruminants, with an additional day with an average temperature > 35°C in the last 12 months increasing the number of dead or lost large ruminants by 40.04%. This high magnitude is explained by the rare occurrence of such days in the sample, with the mean number of days with an average temperature > 35°C being 0.78 per year. Thus, the effect being modelled represents a 128% increase in such days. I find no evidence that these results are driven by increased slaughter, either for sale or own-consumption.<sup>24</sup> This finding, where extreme heat reduces large but not small ruminant herds, is directly supported by the biological literature. Large ruminants, particularly high-production cattle, generate significant internal (metabolic) heat to stay alive, grow, or produce milk. Their large body mass, relative to their skin's surface area, makes it physically difficult for them to shed this internal heat. Small ruminants, in contrast, generate less metabolic heat and have a larger surface-area-to-mass ratio, allowing them to cool down more efficiently and be more resilient to heat shocks (Kadzere et al., 2002; Silanikove, 2000). Finally, I find that the purchase of small ruminants declines by 9.28% for each additional day with an average temperature > 35°C. This suggests that the income shock imposed by extreme heat prevents households from making such types of asset purchases.

## 5.5 Robustness checks

Table 6 presents several checks on the robustness of my main results to alternative model specifications. I only report the estimate associated with HDD, with each row representing a different specification.

Row 1 shows robustness to a less parsimonious model that includes household-level controls, which may be time-varying and thus not captured by household fixed effects, as well as a soil fertility index constructed by Bentze and Wollburg (2024). This index is a composite measure of soil quality, built using principal component analysis to aggregate seven binary

---

<sup>24</sup>Results are available upon request.




Figure 7. **Temperatures and livestock heads under management.** Poisson pseudo-maximum likelihood estimations using the computationally efficient estimator for Poisson regressions using high-dimensional fixed effects developed by [Correia \(2016\)](#). Coefficients are exponentiated and presented as incidence rate ratios (IRR). Coefficients are presented as incidence rate ratios. Reference bin < 31C. Weather over the last 12 months (number of days in each bin). Mean number of annual days with average temperature > 35C: 0.6. 95% confidence intervals are built using robust clustered standard errors at the household level.

indicators of soil constraints.<sup>25</sup> Row 2 estimates the main specification with robust clustered standard errors at the household level instead of spatial HAC standard errors. The statistical significance of my results remains unchanged under this alternative (and arguably simpler) assumption about the error structure. A partial refresh of the panel sample was undertaken in wave 4 ([Table C1](#)). Additionally, the representativeness of the survey was compromised during wave 4 due to security issues, which prevented data collection in certain locations. In row 3, I show the robustness of my findings to the exclusion of the latter waves 4 and 5. These later waves include only a subsample of panel households and rely on a long consumption recall period (30 days). Lastly, in row 4, I allow for different HDD thresholds for the Southern and Northern

<sup>25</sup>The binary indicators are: nutrient availability, nutrient retention, rooting conditions, oxygen availability, excess salts, toxicity, and workability. These indicators are provided with the LSMS-ISA microdata.

	Harvest (kg)	HDDS	Adq prot	Adq iron
1. HH & soil char controls	-0.0154*** (0.0047)	-0.0050* (0.0028)	-0.0019** (0.0008)	-0.0020** (0.0009)
2. Clustered SE at HH level	-0.0154*** (0.0020)	-0.0053** (0.0022)	-0.0022*** (0.0005)	-0.0025*** (0.0006)
3. Excluding waves 4 and 5	-0.0140*** (0.0051)	-0.0069** (0.0032)	-0.0027*** (0.0010)	-0.0029*** (0.0011)
4. Different HDD thr by region	-0.0117** (0.0049)	-0.0055* (0.0029)	-0.0034*** (0.0009)	-0.0044*** (0.0011)

Table 6. **Robustness checks.** This table displays the results from 16 regressions, under the main specification. HH and soil characteristics added: head age, head age squared, head sex, head education, log of HH size, and weighted average soil fertility index (plot level). Regional HDD thresholds: Southern 30°C, Northern 31°C. Adequacy indicators are set to 80% adequacy levels. Total harvest is measured in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional growing season HDD. Coefficients on HDDS can be interpreted as the absolute change in the HDDS for one additional HDD. Coefficients on nutrient adequacy can be interpreted as percentage-point change in the likelihood for one additional HDD. Conley spatial HAC standard errors (in parentheses, except if otherwise stated). Adq: Adequacy. HDDS: Household dietary diversity score. Adq: adequacy. HH: households. prot: protein. SE: standard errors. thr: threshold.

zones.<sup>26</sup> Results are similar to the baseline specification, with a slightly lower coefficient for total harvest and higher coefficients for protein and iron adequacy (in absolute terms).

My identification hinges on the absence of unobserved factors that could plausibly explain the estimated effects. To probe the assumption and verify that high growing season temperatures truly drive the documented effects, I carry out similar falsification exercises as in [Mayorga et al. \(2025\)](#). I randomly reassign temperatures and precipitations across clusters, generate 1,000 such placebo weather datasets, and re-estimate my main specification. Because this permutation breaks the spatial link between actual weather and outcomes, I expect no systematic relationship between heat exposure and changes in the dependent variables. [Figure B13](#) plots the sampling distribution of the placebo estimates of the HDD coefficients, alongside the corresponding point estimate from [Table C3](#), [Table 2](#), and [Figure 5](#). The placebo distributions are approximately

<sup>26</sup>These zone-specific thresholds were chosen by replicating the analysis shown in [Figure 4](#) in the Southern and Northern zones separately. HDD thresholds with the best model fit were respectively 30°C and 31°C. The results from this exercise are available upon request.

normal and centered at zero. In contrast, my main results coefficients (vertical dotted lines) lie far from the tails of the placebo distributions, indicating that the observed effects are unlikely to be due to random assignment.

I also probe how sensitive the results are to alternative choices of the distance and temporal lag parameters used for [Conley \(1999\)](#)'s spatial-HAC standard errors. [Table C11](#) shows that my main results remain stable across distance cutoffs from 50 to 200 km and temporal lags from 1 to 10 years. Using lower distance cut-offs leads to higher statistical significance for the HDDS estimate. However, given the likely significant spatial autocorrelation in both the independent variable and dependent variables, a wider distance cutoff is justified.

In this study, I express the continuous outcomes (mainly energy and nutrient intake) in logarithms because the data are very unevenly distributed (positively skewed) ([Figure B3](#)). Taking the natural log reduces skewness and stabilizes variance. This transformation also yields elasticities in interpretation. I provide a robustness check measuring household energy intake at the food group level in levels rather than logarithms (i.e., in kilocalories). I then divide the effect by the mean intake to express results as a marginal change for one additional HDD in the last growing season. The results of this sensitivity analysis in [Figure B14](#) show that the direction and significance of coefficients are similar to [Figure B10](#).

## **5.6 Additional pathways**

### *Prices as omitted variables*

A potential concern is that my results might be influenced by changes in relative prices. Extreme heat shocks can reduce aggregate supply and increase crop and food prices. A crop price increase may create incentives to increase harvest sales. On the other hand, a price increase in nutritious foods may incentivize reliance on staples and own-produced foods to ensure energy sufficiency, potentially leading to a reduction in commercialization. The main specification includes growing season fixed effects. If agricultural and food markets are well integrated nationally, this approach would control for such price effects. However, this is unlikely to be the case in Nigeria, where evidence points to incomplete or uneven market integration ([Dillon and Barrett, 2016](#); [Amare et al., 2024](#)). Thus, I test the impact of extreme heat in the growing

season on post-harvest community-level (i.e., enumeration area or cluster) food prices using the main specification. Prices are collected during the same survey month as for food expenditure and consumption information in a given enumeration area. I have selected the following food items because of their highest coverage in the data and to represent each of the nine HDDS food groups used in this analysis: maize, potato, onion, banana, beef, egg, fish, groundnut, and milk. Current (nominal) retail prices in local currency (Naira) are harmonised per kilogram and converted to 2015 constant (real) prices using the food consumer price index from the UN Food and Agriculture Organization (FAO) (FAO, 2025).

For the majority of food groups, Table C12 shows that there is no statistically significant relationship between growing season heat and post-harvest local prices. There is one notable exception. An additional HDD in the last growing season increases the price of eggs by 2.96%. The overall lack of a price response is consistent with findings from a previous cross-sectional analysis in Nigeria (Dillon et al., 2015). These results likely reflect my focus on the immediate post-harvest period, whereas price adjustments may take several months to materialize as stocks deplete and markets clear. For instance, evidence from Niger shows that drought-induced price increases emerge only around six months after harvest (Kakpo et al., 2022).

#### *The role of post-harvest off-farm employment*

I consider the effects of extreme heat on *ex-post* labor activity, which could impact income and thus purchasing power. High temperatures can negatively impact employment by either impairing worker health or reducing overall labor demand. Households can also seek additional employment as an adaptation response to compensate for the negative effects of extreme heat on agricultural income (Colmer, 2021). Table 7 shows that point estimates are near zero and that heat in the growing season has no statistically significant effects on post-harvest small-own business or wage employment, either on the extensive or intensive margin. Given the low baseline engagement in wage employment (7.3%) and modest participation in small own-business (18.1%), scope to expand off-farm hours appears limited. This pattern is consistent with binding time and market frictions: households do not substantially reallocate from farm to non-farm work, nor do they add net hours off-farm, even after experiencing adverse heat. These

findings are consistent with thin and incomplete labor markets (Rosenzweig, 1988). The labor-income smoothing margin remains muted, helping explain why expenditure falls (Table 3).

	Extensive margin			Intensive margin			Last 12m
	Farm	SOB	Wage	Farm	SOB	Wage	NFB
GDD in growing season	-0.0002 (0.0001)	0.0000 (0.0000)	0.0000 (0.0000)	-0.0004* (0.0002)	-0.0000 (0.0001)	0.0000 (0.0000)	0.0000 (0.0001)
HDD in growing season	-0.0007 (0.0007)	0.0002 (0.0005)	0.0001 (0.0003)	-0.0018 (0.0017)	-0.0001 (0.0009)	0.0004 (0.0004)	0.0003 (0.0007)
Household FE	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.552	0.539	0.583	0.746	0.865	0.917	0.634
Observations	8,937	8,937	8,931	8,930	8,938	8,938	9,006
Mean Y	0.509	0.181	0.073	.	.	.	0.519

Table 7. **Effect of HDD on post-harvest labor activity.** Labor activity in the last 7 days before interview. Conley spatial HAC standard errors (in parentheses). FE: fixed effects. NFB: Non-farm business. SOB: small-own business.

### *Changes in household composition*

Extreme heat during the growing season may reshape household composition. This could have consequences on the farm’s productive capacity, for example, if it changes the number of working-age household members. It could also change the total amount of food necessary for the household. Three effects may be at play: fertility, migration, and mortality. I test these channels by estimating regressions from my main specification with the number of children, the number of working-age adults, and the number of elderly in the household as the outcome. Results are shown in Table C13. No coefficient is statistically significant at any conventional level. As expected from the climate-health literature, the sign of the coefficient on HDD is negative for young children and the elderly. The latter are more likely to die from heat shocks (Carleton et al., 2022), whereas for the former, fertility may be negatively impacted, for example, by increasing the risk of delivery complications for pregnant women (Barreca and Schaller, 2020). On the other hand, the coefficient on HDD is positive for the number of working-age adults in the household. Interpreted through a liquidity-constraint lens in poor rural settings, hot growing seasons may temporarily inhibit out-migration and induce short-run co-residence or even return

migration (Hirvonen, 2016; Cattaneo and Peri, 2016), increasing the number of working-age adults at home, even in the absence of a higher farm labor demand (Table C15).

#### *Other coping mechanisms*

My main results suggest that farm households respond to temperature-induced crop losses (*ex-post*) by selling a smaller share of harvested crops to maintain calories, which consequently tightens cash and reduces purchases of nutrient-dense foods, thereby degrading dietary quality. In this section, I examine other coping mechanisms that have been previously documented.

First, I examine *ex-ante* productive adaptation responses in terms of input use and crop mix. Consistent with previous studies (Mayorga et al., 2025; Jagnani et al., 2019), I find that households reduce the use of productivity boosting inputs, such as fertilizers. However, I do not find evidence of increased use of pesticides as protective inputs. While mixed cropping, as opposed to monocropping, may allow farmers to dilute their risk from adverse productive shocks (Shaffril et al., 2018), I find no evidence of increased mixed cropping in hotter seasons (Table C14).

Second, I study the effect of growing season heat on farm labor allocation during the agricultural season. Table C15 shows no direct relationship between extreme heat and farm labor inputs. This is consistent with previous results in Nigeria (Mayorga et al., 2025). From a consumption-smoothing perspective, the relevant margin is off-farm earnings. Unfortunately, the LSMS-ISA does not record within-agricultural-season off-farm labor. Prior work in other contexts find that extreme weather increases off-farm labor supply as households seek liquidity (Branco and Féres, 2021; Kochar, 1999). If households face a binding time constraint, holding on-farm work fixed leaves little slack to expand off-farm work, implying a limited scope for within-agricultural-season labor reallocation toward cash income. Lastly, I find no effect of extreme heat in the growing season on the likelihood of a household owning a non-farm business (Table 7).

## 5.7 Back-of-the-envelope uniform warming projections

I provide a projection exercise in which I simulate changes in harvest, dietary diversity, and protein and iron adequacy under a uniform  $+1^{\circ}\text{C}$  warming scenario. This approximately corresponds to the projected warming in Nigeria under the SSP2-4.5 “intermediate” greenhouse gas emissions scenario by 2050 based on the Coupled Model Intercomparison Project Phase 6 (CMIP6) (World Bank, 2024).<sup>27</sup> I calculate the expected change in GDD and HDD and then estimate the expected effect on the outcomes of interest by multiplying the variations by the corresponding estimates from my main specification (Table C3, Table 2, and Figure 5). This exercise deliberately isolates the impact of temperature, employing a *ceteris paribus* assumption. It does not endogenize other complex phenomena associated with climate change, such as biophysical shifts or broader socioeconomic adaptations. Consequently, the findings should not be interpreted as a comprehensive forecast.

Results are reported in Table 8. I find that a uniform  $+1^{\circ}\text{C}$  warming scenario would decrease harvest by 27.38% and subsequently dietary diversity by 0.29%, protein adequacy by 4.04 percentage points (3.18 p.p. for 80% adequacy), and iron adequacy by 1.56 percentage points (3.54 p.p. for 80% adequacy). These percentage changes correspond to an increase of around 1.62 million and 0.63 million households with inadequate protein and iron intake (1.28 and 1.42 million households for 80% adequacy), respectively, out of Nigeria’s 2022 farm household population of 42 million households (National Bureau of Statistics, 2024).

The projected yield impact is larger than the latest  $-10.8\%$  estimate for Africa by 2050 under RCP4.5 from Hultgren et al. (2025). The latter is based on subnational aggregate agricultural production data. Nevertheless, the results from my projection exercise are in line with those of Wollburg et al. (2024), who found that climate shocks reduced crop production among smallholder farmers by 29% in Sub-Saharan Africa between 2008 and 2019, using LSMS-ISA data. While smallholder farmers represent the majority of crop production in Sub-Saharan Africa, the commercial farm sector is rapidly increasing (Jayne et al., 2022). Commercial farming may be

---

<sup>27</sup>Mean annual surface temperature in Nigeria was  $27.7^{\circ}\text{C}$  in 2014 (last year of the historical reference period 1950-2014, used to train the CMIP6) and is expected to be  $28.65^{\circ}\text{C}$  by 2050 under SSP2-4.5 “moderate” scenario by 2050 based on CMIP6 (World Bank, 2024).

Effect of uniform +1°C temperature increase	
Δ GDD	43.601
Δ HDD	12.951
Δ Harvest (%)	-27.378
Δ HDDS (%)	-0.288
Δ Protein 100% adequacy (p.p.)	-4.038
Δ Protein 80% adequacy (p.p.)	-3.183
Δ Iron 100% adequacy (p.p.)	-1.556
Δ Iron 80% adequacy (p.p.)	-3.538

Table 8. **Predicted effects of a uniform +1°C warming scenario.** HDDS: household dietary diversity score. p.p.: percentage points.

more resilient to climate shocks, with larger farm sizes and better access to agricultural inputs.

## 6 Discussion

This paper set out to identify how extreme heat during the growing season affects post-harvest nutrition outcomes for smallholder farm households in Nigeria, extending beyond energy sufficiency to focus on nutrient adequacy. A central objective was to shed light on household adaptation responses to temperature-induced crop losses in terms of consumption, commercialization, and purchases. Using panel microdata and exploiting quasi-random within-household variations in weather across growing seasons, I demonstrate that extreme heat reduces harvested quantities but has no immediate impact on post-harvest energy intake, while exacerbating micronutrient shortfalls.

I document a behavioral response to the production shock. Households increase reliance on own-produced foods and retain a larger share of their harvest, mainly composed of cereals and tubers, to preserve calorie intake. Combined with the temperature-induced harvest shortfall, the decreased share of harvested crops sold reduces cash income and food purchases, with cuts falling disproportionately on cash-intensive and nutrient-dense items, such as vegetables, pulses, and meat. While safeguarding energy intake, this *ex-post* adaptation strategy degrades diet quality. This is consistent with predictions from non-separable farm household models in

the presence of incomplete food and labor markets and binding caloric constraints.

A uniform  $+1^{\circ}\text{C}$  warming is associated with an additional 1.62 million and 0.63 million Nigerian households with inadequate protein and iron intake. The nutritional impacts are significantly larger for households with children under five years old, a concerning finding given the crucial role of these nutrients in linear growth, cognitive development, and immune function during early childhood (Black et al., 2013). Moreover, these estimates are likely lower bounds. Climate change may further erode dietary quality by lowering the nutrient content of foods, thereby amplifying nutrient inadequacies (IPCC, 2019). Although my estimates capture only responses up to four months post-harvest, I show that adverse nutritional effects worsen as time since harvest elapses. As stocks deplete, the ex-post strategy that initially safeguards calories may also fail to maintain energy sufficiency; a transition that future work could track with high-frequency panel survey data.

To my knowledge, Stainier et al. (2025) represents the only other study that separates own-produced from purchased foods when mapping climate shocks to diets. Both studies document diet-quality losses following hot growing seasons; however, the direction of the market response differs. In rural India, Stainier et al. (2025) find households purchase more to offset own-production shortfalls, enabled by a labor reallocation into non-agricultural work. In Nigeria, I find no post-harvest expansion of off-farm labor and, consequently, no compensating rise in food purchases; instead, households sell less and buy less. This may be explained by relatively more complete rural labor markets in India.<sup>28</sup> The absence of an operative labor channel in Nigeria maps into a defensive autarkic behavior and opposite purchase responses.

I acknowledge a number of limitations to my analysis. First, food consumption is recorded with a 7-day recall in the first three waves and a 30-day recall in the last two. Shorter windows tend to overstate mean intake (Mukherjee and Chaudhury, 2020). While this primarily affects levels, bias could arise if recall behavior covaries with heat. Nevertheless, Villacis et al. (2023) show that both recall periods perform as well in identifying food-insecure households in Nigeria. I mitigate this by including growing-season (survey-wave) fixed effects that

---

<sup>28</sup>Also, Stainier et al. (2025) do not examine the impact on commercialization and include rural non-farm households in their sample.

absorb recall-window differences and common shocks. I also show the robustness of my results to excluding the last two waves. Second, because food may not be distributed equally within households (D'Souza and Tandon, 2019), the effects of extreme heat on undernutrition may be attenuated or masked for vulnerable subgroups, particularly if allocation shifts under financial stress (Hazrana et al., 2025). I partly address this by examining heterogeneity by household composition. However, a more comprehensive assessment would require individual-level consumption data. Lastly, although my analysis accounts for zone-specific growing season calendars and degree days thresholds (i.e., Southern vs. Northern), a more granular approach, tailoring season timing and temperature cut-offs to crop-specific phenology and local climate, would likely improve precision. Future work with finer seasonal data could reduce timing mismatches and strengthen inference across heterogeneous agro-climatic contexts.

Despite these shortcomings, this paper makes significant contributions to our understanding of the consequences of extreme temperatures for household food utilization and nutrition, an issue that remains understudied in the literature. Shifting from coarse indices to nutrient-specific intake and adequacy measures reveals declines in diet quality that were previously obscured by aggregate calorie intake metrics. Designing climate adaptation policy requires understanding household responses to heat. Crucially, all core variables in this analysis (agricultural production, livestock management, prices, consumption, and nutrition) are measured within the same household-panel architecture, improving the internal consistency of estimated pathways and reducing the scope for ecological fallacy.

Prior work shows that commercialization generally improves diet quality (Chegere and Kauky, 2022; Ogutu et al., 2020), yet its nutrition gains are attenuated in the presence of climate shocks (Hazrana et al., 2025). However, the behavioral channel through which weather shocks attenuate these gains has remained underexplored. By separating own-produced from purchased intake and investigating the commercialization–consumption trade-off, this paper reveals a margin of adjustment not previously documented: households reduce the share of harvest sold and secure energy intake through home-grown staple consumption, while cutting purchases of nutrient-dense foods. Thus, climate shocks not only dampen the returns to commercialization but also reduce commercialization itself, tightening cash constraints and transmitting the shock

into diet-quality losses. This immediate post-harvest response erodes the commercialization pathway out of poverty ([Barrett, 2008](#)) and is likely relevant in other smallholder contexts with incomplete markets.

Safeguarding nutrition under climate change necessitates moving beyond caloric sufficiency to target nutrient-adequate diets that support healthy and active lives. It requires integrating climate-risk mitigation with measures that keep smallholders engaged in markets and improve their access to non-farm labor opportunities. Direct financial support could also help mitigate nutritional losses ([Premand and Stoeffler, 2022](#)).

## References

- Akombi, Blessing Jaka, Kingsley Emwinyore Agho, John Joseph Hall, Dafna Merom, Thomas Astell-Burt, and Andre M. N. Renzaho**, “Stunting and severe stunting among children under-5 years in Nigeria: A multi-level analysis,” *BMC Pediatrics*, 2017, 17 (1), 15.
- Amare, Mulubrhan, Bekele Shiferaw, Hiroyuki Takeshima, and George Mavrotas**, “Variability in agricultural productivity and rural household consumption inequality: Evidence from Nigeria and Uganda,” *Agricultural Economics*, 2021, 52 (1), 19–36.
- , **Kibrom A. Abay, and Patrick Hatzenbuehler**, “Spatial market integration during a pandemic: Evidence from food markets in Nigeria,” *Agricultural Economics*, 2024, 55 (1), 86–103.
- Aragón, Fernando M., Francisco Oteiza, and Juan Pablo Rud**, “Climate Change and Agriculture: Subsistence Farmers’ Response to Extreme Heat,” *American Economic Journal: Economic Policy*, February 2021, 13 (1), 1–35.
- Babatunde, Raphael O. and Matin Qaim**, “Impact of off-farm income on food security and nutrition in Nigeria,” *Food Policy*, 2010, 35 (4), 303–311.
- Bai, Yan, Robel Alemu, Steven A. Block, Derek Headey, and William A. Masters**, “Cost and affordability of nutritious diets at retail prices: Evidence from 177 countries,” *Food Policy*, 2021, 99, 101983.
- Barreca, Alan and Jessamyn Schaller**, “The impact of high ambient temperatures on delivery timing and gestational lengths,” *Nature Climate Change*, 2020, 10 (1), 77–82.
- Barrett, Christopher B.**, “Smallholder market participation: Concepts and evidence from eastern and southern Africa,” *Food Policy*, 2008, 33 (4), 299–317.
- Benjamin, Dwayne**, “Household Composition, Labor Markets, and Labor Demand: Testing for Separation in Agricultural Household Models,” *Econometrica*, 1992, 60 (2), 287–322.
- Bentze, Thomas and Philip Wollburg**, “A longitudinal cross-country dataset on agricultural productivity and welfare in Sub-Saharan Africa,” Working Paper 10976, World Bank November 2024.
- Black, Robert E, Cesar G Victora, Susan P Walker, Zulfiqar A Bhutta, Parul Christian, Mercedes de Onis, Majid Ezzati, Sally Grantham-McGregor, Joanne Katz, Reynaldo Martorell, and Ricardo Uauy**, “Maternal and child undernutrition and overweight in low-income and middle-income countries,” *The Lancet*, 2013, 382 (9890), 427–451.
- Blakeslee, David S. and Ram Fishman**, “Weather Shocks, Agriculture, and Crime,” *Journal of Human Resources*, 2018, 53 (3), 750–782.
- Blom, Sylvia, Ariel Ortiz-Bobea, and John Hoddinott**, “Heat exposure and child nutrition: Evidence from West Africa,” *Journal of Environmental Economics and Management*, 2022, 115, 102698.
- Branco, Danyelle and José Féres**, “Weather Shocks and Labor Allocation: Evidence from Rural Brazil,” *American Journal of Agricultural Economics*, 2021, 103 (4), 1359–1377.
- Burke, Marshall and Kyle Emerick**, “Adaptation to Climate Change: Evidence from US Agriculture,” *American Economic Journal: Economic Policy*, August 2016, 8 (3), 106–40.
- Carleton, Tamma, Amir Jina, Michael Delgado, Michael Greenstone, Trevor Houser, Solomon Hsiang, Andrew Hultgren, Robert E Kopp, Kelly E McCusker, Ishan Nath, James Rising, Ashwin Rode, Hee Kwon Seo, Arvid Viaene, Jiacan Yuan, and Alice Tianbo Zhang**, “Valuing the Global Mortality Consequences of Climate Change Accounting for Adaptation Costs and Benefits\*,” *The Quarterly Journal of Economics*, 04 2022, 137 (4), 2037–2105.
- , **Esther Duflo, B. Kelsey Jack, and Guglielmo Zappalà**, “Chapter 4 - Adaptation to climate change,” in Lint Barrage and Solomon Hsiang, eds., *Lint Barrage and Solomon Hsiang, eds.*, Vol. 1 of *Handbook of the Economics of Climate Change*, North-Holland, 2024, pp. 143–248.
- Carletto, Gero, Marco Tiberti, and Alberto Zezza**, “Measure for Measure: Comparing Survey Based Estimates of Income and Consumption for Rural Households,” *The World Bank Research Observer*, 06 2021, 37 (1), 1–38.
- Carpenna, Fenella**, “How do droughts impact household food consumption and nutritional intake? A study of rural India,” *World Development*, 2019, 122, 349–369.
- Case, Anne and Christina Paxson**, “Stature and Status: Height, Ability, and Labor Market Outcomes,” *Journal of Political Economy*, 2008, 116 (3), 499–532.
- Cattaneo, Cristina and Giovanni Peri**, “The migration response to increasing temperatures,” *Journal of Development Economics*, 2016, 122, 127–146.
- Chegere, Martin Julius and Monica Sebastian Kauky**, “Agriculture commercialisation, household dietary di-

- versity and nutrition in Tanzania,” *Food Policy*, 2022, 113, 102341. Role of Policy in Reducing Malnutrition in sub-Saharan Africa.
- Colen, L., P.C. Melo, Y. Abdul-Salam, D. Roberts, S. Mary, and S. Gomez Y Paloma**, “Income elasticities for food, calories and nutrients across Africa: A meta-analysis,” *Food Policy*, 2018, 77, 116–132.
- Colmer, Jonathan**, “Temperature, Labor Reallocation, and Industrial Production: Evidence from India,” *American Economic Journal: Applied Economics*, October 2021, 13 (4), 101–24.
- Conley, T.G.**, “GMM estimation with cross sectional dependence,” *Journal of Econometrics*, 1999, 92 (1), 1–45.
- Correia, Sergio**, “Linear Models with High-Dimensional Fixed Effects: An Efficient and Feasible Estimator,” Working Paper 2016.
- Costinot, Arnaud, Dave Donaldson, and Cory Smith**, “Evolving Comparative Advantage and the Impact of Climate Change in Agricultural Markets: Evidence from 1.7 Million Fields around the World,” *Journal of Political Economy*, 2016, 124 (1), 205–248.
- Dasgupta, Shouro and Elizabeth J. Z. Robinson**, “Attributing changes in food insecurity to a changing climate,” *Scientific Reports*, 2022, 12 (1), 4709.
- de Janvry, Alain, Marcel Fafchamps, and Elisabeth Sadoulet**, “Peasant Household Behaviour with Missing Markets: Some Paradoxes Explained,” *The Economic Journal*, 11 1991, 101 (409), 1400–1417.
- Deaton, Angus**, *The Analysis of Household Surveys: A Microeconomic Approach to Development Policy*, Baltimore, MD; Washington, DC: Johns Hopkins University Press for the World Bank, 1997. Published for the World Bank.
- Dell, Melissa, Benjamin F. Jones, and Benjamin A. Olken**, “What Do We Learn from the Weather? The New Climate-Economy Literature,” *Journal of Economic Literature*, September 2014, 52 (3), 740–98.
- Dillon, Andrew, Kevin McGee, and Gbemisola Oseni**, “Agricultural Production, Dietary Diversity and Climate Variability,” *The Journal of Development Studies*, 2015, 51 (8), 976–995.
- Dillon, Brian and Christopher B. Barrett**, “Agricultural factor markets in Sub-Saharan Africa: An updated view with formal tests for market failure,” *Food Policy*, 2017, 67, 64–77. Agriculture in Africa – Telling Myths from Facts.
- Dillon, Brian M. and Christopher B. Barrett**, “Global Oil Prices and Local Food Prices: Evidence from East Africa,” *American Journal of Agricultural Economics*, 2016, 98 (1), 154–171.
- Dillon, Brian, Peter Brummund, and Germano Mwabu**, “Asymmetric non-separation and rural labor markets,” *Journal of Development Economics*, 2019, 139, 78–96.
- D’Souza, Anna and Sharad Tandon**, “Intrahousehold Nutritional Inequities in Rural Bangladesh,” *Economic Development and Cultural Change*, 2019, 67 (3), 625–657.
- Ecker, Olivier and Patrick L. Hatzenbuehler**, “Food consumption–production response to agricultural policy and macroeconomic change in Nigeria,” *Applied Economic Perspectives and Policy*, 2022, 44 (2), 982–1002.
- Falco, Salvatore Di, Marcella Veronesi, and Mahmud Yesuf**, “Does Adaptation to Climate Change Provide Food Security? A Micro-Perspective from Ethiopia,” *American Journal of Agricultural Economics*, 2011, 93 (3), 829–846.
- FAO**, “Guidelines for measuring household and individual dietary diversity,” Report, Food and Agricultural Organization of the United Nations 2011.
- FAO**, *The State of Food and Agriculture 2013: Food systems for better nutrition*, Rome: Food and Agriculture Organization of the United Nations, 2013.
- FAO**, “FAOSTAT: Undernourishment — Sub-Saharan Africa, 2022,” <https://www.fao.org/faostat/en/2022>. Dataset. Accessed 2025-11-01.
- , “FAOSTAT: Consumer Prices,” <https://www.fao.org/faostat/en/#data/CP> 2025. Accessed 2025-11-07.
- FAO, UN ECA, and AUC**, *Africa – Regional Overview of Food Security and Nutrition 2021: Statistics and trends*, Accra: Food and Agriculture Organization of the United Nations, 2021.
- Funk, Chris, Pete Peterson, Martin Landsfeld, Diego Pedreros, James Verdin, Shraddhanand Shukla, Gregory Husak, James Rowland, Laura Harrison, Andrew Hoell, and Joel Michaelsen**, “The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes,” *Scientific Data*, 2015, 2 (150066), 2052–4463.
- Glewwe, Paul, Hanan G Jacoby, and Elizabeth M King**, “Early childhood nutrition and academic achievement: a longitudinal analysis,” *Journal of Public Economics*, 2001, 81 (3), 345–368.

- Hazrana, Jaweriah and Ashok K. Mishra**, “Commercialisation, exogenous shocks and nutrition: Evidence from smallholder farmers in Bangladesh,” *Journal of Agricultural Economics*, 2025, 76 (1), 98–138.
- , **Pratap S. Birthal, and Ashok K. Mishra**, “Equal exposure, unequal effects of climate change: Gendered impacts on food consumption and nutrition in rural Bangladesh,” *Ecological Economics*, 2025, 230, 108486.
- Headey, Derek, Kalle Hirvonen, and John Hoddinott**, “Animal Sourced Foods and Child Stunting,” *American Journal of Agricultural Economics*, 2018, 100 (5), 1302–1319.
- , **Olivier Ecker, and Jean-Francois Trinh Tan**, *Shocks to the system: monitoring food security in a volatile world*, Cheltenham, UK; Northampton, MA: Edward Elgar Publishing, March 2014.
- Hersbach, Hans, Bill Bell, Paul Berrisford, Shoji Hirahara, András Horányi, Joaquín Muñoz-Sabater, Julien Nicolas, Carole Peubey, Raluca Radu, Dinand Schepers, Adrian Simmons, Cornel Soci, Saleh Abdalla, Xavier Abellan, Gianpaolo Balsamo, Peter Bechtold, Gionata Biavati, Jean Bidlot, Massimo Bonavita, Giovanna De Chiara, Per Dahlgren, Dick Dee, Michail Diamantakis, Rossana Dragani, Johannes Flemming, Richard Forbes, Manuel Fuentes, Alan Geer, Leo Haimberger, Sean Healy, Robin J. Hogan, Elías Hólm, Marta Janisková, Sarah Keeley, Patrick Laloyaux, Philippe Lopez, Cristina Lupu, Gabor Radnoti, Patricia de Rosnay, Iryna Rozum, Freja Vamborg, Sebastien Villaume, and Jean-Noël Thépaut**, “The ERA5 global reanalysis,” *Quarterly Journal of the Royal Meteorological Society*, 2020, 146 (730), 1999–2049.
- Hirvonen, Kalle**, “Temperature Changes, Household Consumption, and Internal Migration: Evidence from Tanzania,” *American Journal of Agricultural Economics*, 2016, 98 (4), 1230–1249.
- Hoynes, Hilary, Diane Whitmore Schanzenbach, and Douglas Almond**, “Long-Run Impacts of Childhood Access to the Safety Net,” *American Economic Review*, April 2016, 106 (4), 903–34.
- Hultgren, Andrew, Tamma Carleton, Michael Delgado, Diana R. Gergel, Michael Greenstone, Trevor Houser, Solomon Hsiang, Amir Jina, Robert E. Kopp, Steven B. Malevich, Kelly E. McCusker, Terin Mayer, Ishan Nath, James Rising, Ashwin Rode, and Jiacan Yuan**, “Impacts of climate change on global agriculture accounting for adaptation,” *Nature*, 2025, 642, 644–652.
- IPCC**, *Chapter 5: Food Security*, Geneva: Intergovernmental Panel on Climate Change, 2019.
- Jagnani, Maulik, Christopher B. Barrett, Yanyan Liu, and Liangzhi You**, “Within-Season Response to Warmer Temperatures: Defensive Investments by Kenyan Farmers,” Working Paper 314, African Development Bank Group, Abidjan, Côte d’Ivoire July 2019.
- Jayne, T.S., Ayala Wineman, Jordan Chamberlin, Milu Muyanga, and Felix Kwame Yeboah**, “Changing Farm Size Distributions and Agricultural Transformation in Sub-Saharan Africa,” *Annual Review of Resource Economics*, 2022, 14 (Volume 14, 2022), 109–130.
- Kadzere, C.T, M.R Murphy, N Silanikove, and E Maltz**, “Heat stress in lactating dairy cows: a review,” *Livestock Production Science*, 2002, 77 (1), 59–91.
- Kakpo, Ange, Bradford F. Mills, and Stéphanie Brunelin**, “Weather shocks and food price seasonality in Sub-Saharan Africa: Evidence from Niger,” *Food Policy*, 2022, 112, 102347.
- Kocher, Anjini**, “Smoothing Consumption by Smoothing Income: Hours-of-Work Responses to Idiosyncratic Agricultural Shocks in Rural India,” *The Review of Economics and Statistics*, 02 1999, 81 (1), 50–61.
- Kroeger, Carolin**, “Heat is associated with short-term increases in household food insecurity in 150 countries and this is mediated by income,” *Nature Human Behaviour*, 2023, 7 (10), 1777–1786.
- Malacarne, J.G. and L.A. Paul**, “Do the benefits of improved management practices to nutritional outcomes “dry up” in the presence of drought? Evidence from East Africa,” *Food Policy*, 2022, 113, 102332. Role of Policy in Reducing Malnutrition in sub-Saharan Africa.
- Maluccio, John A., John Hoddinott, Jere R. Behrman, Reynaldo Martorell, Agnes R. Quisumbing, and Aryeh D. Stein**, “The Impact of Improving Nutrition During Early Childhood on Education among Guatemalan Adults,” *The Economic Journal*, 03 2009, 119 (537), 734–763.
- Masters, William A., Yan Bai, Anna Herforth, Daniel B. Sarpong, Fulgence Mishili, Joyce Kinabo, and Jennifer C. Coates**, “Measuring the Affordability of Nutritious Diets in Africa: Price Indexes for Diet Diversity and the Cost of Nutrient Adequacy,” *American Journal of Agricultural Economics*, 2018, 100 (5), 1285–1301.
- Mayorga, Joaquin, Alexis H Villacis, and Ashok K Mishra**, “Farm-level agricultural productivity and adaptation to extreme heat,” *American Journal of Agricultural Economics*, 2025, pp. 1–23.
- McCullough, Ellen B., Meichen Lu, Yawotse Nouve, Joanne Arsenaault, and Chen Zhen**, “Nutrient adequacy for poor households in Africa would improve with higher income but not necessarily with lower food prices,” *Nature Food*, 2024, 5, 171–181.

- Mukherjee, Diganta and Prabir Chaudhury**, “Why does “last week” reporting give higher estimates than “last month”?” *Communications in Statistics - Theory and Methods*, 2020, 49 (8), 1873–1893.
- National Bureau of Statistics**, “National Agricultural Sample Census (NASC) Report 2022,” 2024. Launched June 28, 2024.
- Nguyen, Thanh-Tung and Matin Qaim**, “Local and regional food production diversity are positively associated with household dietary diversity in rural Africa,” *Nature Food*, 2025, 6 (2), 205–212.
- Ogotu, Sylvester Ochieng, Theda Gödecke, and Matin Qaim**, “Agricultural Commercialisation and Nutrition in Smallholder Farm Households,” *Journal of Agricultural Economics*, 2020, 71 (2), 534–555.
- Omiti, John M., David J. Otieno, Timothy O. Nyanamba, and Ellen McCullough**, “Factors Influencing the Intensity of Market Participation by Smallholder Farmers: A Case Study of Rural and Peri-Urban Areas of Kenya,” *African Journal of Agricultural and Resource Economics*, 2009, 3 (1), 57–82.
- Passarelli, Simone, Christopher M Free, Alon Shepon, Ty Beal, Carolina Batis, and Christopher D Golden**, “Global estimation of dietary micronutrient inadequacies: a modelling analysis,” *The Lancet Global Health*, 2024, 12 (10), e1590–e1599.
- Pitt, Mark M. and Mark R. Rosenzweig**, “Health and Nutrient Consumption Across and Within Farm Households,” *The Review of Economics and Statistics*, 1985, 67 (2), 212–223.
- Premand, Patrick and Quentin Stoeffler**, “Cash transfers, climatic shocks and resilience in the Sahel,” *Journal of Environmental Economics and Management*, 2022, 116, 102744.
- Randell, Heather, Clark Gray, and Elizabeth H. Shayo**, “Climatic conditions and household food security: Evidence from Tanzania,” *Food Policy*, 2022, 112, 102362.
- Rashid, Dewan Arif, Lisa C. Smith, and Tauhidur Rahman**, “Determinants of Dietary Quality: Evidence from Bangladesh,” *World Development*, 2011, 39 (12), 2221–2231.
- Rosenzweig, Mark R.**, *Chapter 15 Labor markets in low-income countries*, Vol. 1 of *Handbook of Development Economics*, Elsevier, 1988.
- **and Kenneth I. Wolpin**, “Credit Market Constraints, Consumption Smoothing, and the Accumulation of Durable Production Assets in Low-Income Countries: Investments in Bullocks in India,” *Journal of Political Economy*, 1993, 101 (2), 223–244.
- **and Oded Stark**, “Consumption Smoothing, Migration, and Marriage: Evidence from Rural India,” *Journal of Political Economy*, 1989, 97 (4), 905–926.
- Schlenker, Wolfram and Michael J. Roberts**, “Nonlinear temperature effects indicate severe damages to U.S. crop yields under climate change,” *Proceedings of the National Academy of Sciences*, 2009, 106 (37), 15594–15598.
- **, W. Michael Hanemann, and Anthony C. Fisher**, “The Impact of Global Warming on U.S. Agriculture: An Econometric Analysis of Optimal Growing Conditions,” *The Review of Economics and Statistics*, 02 2006, 88 (1), 113–125.
- Shaffril, Hayrol Azril Mohamed, Steven Eric Krauss, and Samsul Farid Samsuddin**, “A systematic review on Asian’s farmers’ adaptation practices towards climate change,” *Science of The Total Environment*, 2018, 644, 683–695.
- Silanikove, Nissim**, “Effects of heat stress on the welfare of extensively managed domestic ruminants,” *Livestock Production Science*, 2000, 67 (1), 1–18.
- Stainier, Paul, Manisha Shah, and Alan Barreca**, “Hot Weather, Undernutrition, and Adaptation in Rural India,” Working Paper 34047, National Bureau of Economic Research July 2025.
- Verger, Eric O., Terri J. Ballard, Marie Claude Dop, and Yves Martin-Prevel**, “Systematic review of use and interpretation of dietary diversity indicators in nutrition-sensitive agriculture literature,” *Global Food Security*, 2019, 20, 156–169.
- Villacis, Alexis H., Joaquin Mayorga, and Ashok K. Mishra**, “Experience-based food insecurity and agricultural productivity in Nigeria,” *Food Policy*, 2022, 113, 102286. Role of Policy in Reducing Malnutrition in sub-Saharan Africa.
- **, Syed Badruddoza, Ashok K. Mishra, and Joaquin Mayorga**, “The role of recall periods when predicting food insecurity: A machine learning application in Nigeria,” *Global Food Security*, 2023, 36, 100671.
- Wegenast, Tim, Niklas Hänze, Roos Haer, and Marcel Birulés**, “The effect of traditional agricultural practices on the food consumption of households facing extreme weather events in Tanzania,” *Agriculture & Food Security*, January 2025, 13 (1), 56.

- WMO**, “Africa faces disproportionate burden from climate change and adaptation costs,” 2024. Press release. Accessed 2025-11-02.
- Wollburg, Philip, Yannick Markhof, Thomas Bentze, and Giulia Ponzini**, “Substantial impacts of climate shocks in African smallholder agriculture,” *Nature Sustainability*, 2024, 7, 1525–1534.
- World Bank**, “Nigeria—Agriculture and Rural Poverty: A Policy Note,” Technical Report Report No. 78364-NG, World Bank, Washington, DC 2014. Poverty Reduction and Economic Management, Africa Region.
- , “Climate Risk Country Profile: Nigeria,” Technical Report, World Bank Group, Washington, DC 2021. “Please cite the work as follows: *Climate Risk Country Profile: Nigeria* (2021): The World Bank Group.”
- , “Nigeria — Climate Data & Projections (Climate Change Knowledge Portal),” 2024. Climate Change Knowledge Portal.
- , “The World Bank in Nigeria: Overview,” 2025. Last updated: 2025-10-13.
- World Food Programme and Food and Agriculture Organization of the United Nations**, “Hunger Hotspots. FAO–WFP early warnings on acute food insecurity: June to October 2025 outlook,” Report, WFP and FAO, Rome 2025. CC BY 4.0.

## Appendix

### A. Appendix Data Details

#### LSMS-ISA geographical offset

In the LSMS-ISA data, household confidentiality was preserved by applying a geographic offset process to the location data. Initially, coordinates were aggregated to the enumeration area (EA) level by calculating the mean location of each EA. These aggregate points were then randomly displaced by up to 2 km in urban areas and up to 5 km in rural areas, with the latter receiving a larger offset due to a higher disclosure risk in less dense settings. To further mask the data, a small subset of rural clusters (1% in most samples) was subjected to a larger displacement of up to 10 km. The maximum potential offset for any rural point is 10 km.

#### Weather data

I extract minimum and maximum daily temperatures from the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 climate reanalysis dataset at the  $0.1^\circ$  resolution, or about  $10 \times 10$  km close to the equator (Hersbach et al., 2020). I estimate the average daily temperature as the average of the daily maximum and daily minimum temperatures. Total precipitation data are extracted from the Climate Hazards Group InfraRed Precipitation with Station (CHIRPS) database, which combines observations from real-time meteorological stations with infra-red data at the  $0.05^\circ$  resolution, or about  $5 \times 5$  km close to the equator (Funk et al., 2015). Total precipitation data is reaggregated using means to match ERA5's resolution. The weather data for a specific  $0.1^\circ$  resolution grid square is assigned to an LSMS-ISA cluster if the cluster lies within that grid square. The low resolution of the weather data ( $\approx 10 \times 10$  km) is larger than the random geographic offsets applied to the coordinates. Therefore, the process of altering the coordinates is not expected to meaningfully affect the effect of the weather variables on the outcome variables.

#### Agricultural output data

Agricultural output data initially provided by the LSMS-ISA surveys at the plot level are aggregated, i.e., summed up, at the household level, while the maximum values of indicator variables (e.g., intercropped or not) are retained. Plots for which output amounts or size (area) are missing or not strictly positive are dropped. Agricultural output values are winsorised at the 99th percentile. Harvest is valued using median prices per enumeration area. If there are fewer than 10 observed sales in the area, prices are calculated at a higher geographical level. Prices are calculated independently for each crop type (Bentze and Wollburg, 2024). Harvest values are converted to current USD and deflated to 2020 USD, using exchange rates and a deflator from the World Bank. The same transformation is performed for household expenditure.

#### Food and nutrient consumption data

Households report food items consumed at home over the seven full days preceding the interview date. Food consumption is categorized into three types: purchased, own-produced, and

gifted (including transfers). Food items are grouped according to the Household Dietary Diversity Score (HDDS) food grouping (FAO, 2011), including: A. Cereals; B. Roots and tubers; C. Vegetables; D. Fruits; E. Meat, poultry, offal; F. Eggs; G. Fish and seafood; H. Pulses, legumes, nuts; I. Milk and milk products; J. Oil/fats; K. Sugar/honey; and L. Miscellaneous.

Food and nutrient consumption data cleaning follows McCullough et al. (2024). Units of consumption are harmonised to kilograms (kg) using conversion factors provided with the survey microdata. The interquartile range method is used to clean outliers of consumption per adult equivalent at the household-item level for each country. The nutritional content of food items is determined using published food composition tables from Africa (Lukmanji et al., 2008; Vincent et al., 2020; Stadlmayr et al., 2012; Hotz et al., 2012; University of California, Berkeley, 2006), and completed with data from the United States Department of Agriculture (USDA) (U.S. Department of Agriculture, Agricultural Research Service, 2019). For staple foods, I incorporate assumed fortification rates based on data from the Global Fortification Data Exchange (Global Fortification Data Exchange, 2023). Further adjustments were made to account for edible portions and nutrient losses during cooking, using information from the USDA (Matthews and Garrison, 1975; U.S. Department of Agriculture, Agricultural Research Service, 2007). As in McCullough et al. (2024), food away from home consumption is not included due to a lack of information about the items consumed to match them with nutrient content information.

I also follow McCullough et al. (2024) in constructing household-level estimated average requirements (EAR) for energy and each nutrient. EAR per adult equivalent is based on the age and sex composition of each household, assuming adults are of average weight and engage in moderate activity levels. Dietary energy (DE) and protein requirements are taken from FAO/WHO/UNU (2004). I establish EAR for vitamin A, total folate, and iron using the work of the Institute of Medicine (2006), for zinc based on the International Zinc Nutrition Consultative Group et al. (2004), and for iron from the Institute of Medicine (2002). I assume low bioavailability for both zinc and iron. For zinc, this is due to diets relying heavily on unrefined cereals, and for iron, it is because diets are high in phytate and low in animal-sourced foods.

To measure adequacy, I calculate the ratio of a household's intake to its EAR. A ratio greater than one indicates sufficient intake, while a ratio lower than one shows the fraction of the EAR being consumed. I then create a binary variable for energy and each nutrient that equals one if a household's intake exceeds its EAR and zero otherwise.

## References for Appendix A

- Bentze, Thomas and Philip Wollburg**, “A longitudinal cross-country dataset on agricultural productivity and welfare in Sub-Saharan Africa,” Working Paper 10976, World Bank November 2024.
- FAO**, “Guidelines for measuring household and individual dietary diversity,” Report, Food and Agricultural Organization of the United Nations 2011.
- FAO/WHO/UNU**, “Human Energy Requirements: Report of a Joint FAO/WHO/UNU Expert Consultation,” Technical Report, Food and Agriculture Organization of the United Nations (FAO); World Health Organization (WHO); United Nations University (UNU), Rome, Italy 2004.
- Funk, Chris, Pete Peterson, Martin Landsfeld, Diego Pedreros, James Verdin, Shraddhanand Shukla, Gregory Husak, James Rowland, Laura Harrison, Andrew Hoell, and Joel Michaelsen**, “The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes,” *Scientific Data*, 2015, 2 (150066), 2052–4463.
- Global Fortification Data Exchange**, “Global Fortification Data Exchange,” 2023.
- Hersbach, Hans, Bill Bell, Paul Berrisford, Shoji Hirahara, András Horányi, Joaquín Muñoz-Sabater, Julien Nicolas, Carole Peubey, Raluca Radu, Dinand Schepers, Adrian Simmons, Cornel Soci, Saleh Abdalla, Xavier Abellan, Gianpaolo Balsamo, Peter Bechtold, Gionata Biavati, Jean Bidlot, Massimo Bonavita, Giovanna De Chiara, Per Dahlgren, Dick Dee, Michail Diamantakis, Rossana Dragani, Johannes Flemming, Richard Forbes, Manuel Fuentes, Alan Geer, Leo Haimberger, Sean Healy, Robin J. Hogan, Elías Hólm, Marta Janisková, Sarah Keeley, Patrick Laloyaux, Philippe Lopez, Cristina Lupu, Gabor Radnoti, Patricia de Rosnay, Iryna Rozum, Freja Vamborg, Sebastien Villaume, and Jean-Noël Thépaut**, “The ERA5 global reanalysis,” *Quarterly Journal of the Royal Meteorological Society*, 2020, 146 (730), 1999–2049.
- Hotz, C., A. Lubowa, C. Sison, M. Moursi, and C. Loechl**, “A Food Composition Table for Central and Eastern Uganda,” HarvestPlus Technical Monograph, International Food Policy Research Institute (IFPRI) and International Center for Tropical Agriculture (CIAT), Washington, DC and Cali, Colombia 2012.
- Institute of Medicine**, *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*, Washington, DC: National Academies Press, 2002.
- , *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*, Washington, DC: National Academies Press, 2006.
- International Zinc Nutrition Consultative Group, K. Brown, J. Rivera, Z. Bhutta, R. Gibson, J. King, B. Lönnerdal, M. Ruel, B. Sandström, E. Wasantwisut, and C. Hotz**, “International Zinc Nutrition Consultative Group (IZiNCG) Technical Document No. 1: Assessment of the Risk of Zinc Deficiency in Populations and Options for Its Control,” *Food and Nutrition Bulletin*, 2004, 25 (1), 99–203.
- Lukmanji, Z., E. Hertzmark, N. Mlingi, V. Assey, G. Ndossi, and W. Fawzi**, *Tanzania Food Composition Tables*, Dar es Salaam, Tanzania: MUHAS; Tanzania Food and Nutrition Centre (TFNC); Harvard School of Public Health (HSPH), 2008. PDF (Sept 2023 update).
- Matthews, R. H. and Y. J. Garrison**, *Food Yields Summarized by Different Stages of Preparation* number 102. In ‘Agriculture Handbook.’, Beltsville, MD: U.S. Department of Agriculture, 1975.
- McCullough, Ellen B., Meichen Lu, Yawotse Nouve, Joanne Arsenault, and Chen Zhen**, “Nutrient adequacy for poor households in Africa would improve with higher income but not necessarily with lower food prices,” *Nature Food*, 2024, 5, 171–181.
- Stadlmayr, B., U. R. Charrondièrre, V. Enujiugha, R. Bayili, E. Fagbohoun, B. Samb, P. Addy, I. Barikmo, F. Ouattara, A. Oshaug, I. Akinyele, G. Annor, K. Bomfeh, H. Ene-Obong, I. Smith, I. Thiam, and B. Burlingame**, *West African Food Composition*, Rome, Italy: Food and Agriculture Organization of the United Nations (FAO), 2012.
- University of California, Berkeley**, “International Minilist: WorldFood Dietary Assessment System, 2nd edition,” Internet 2006.
- U.S. Department of Agriculture, Agricultural Research Service**, “USDA Table of Nutrient Retention Factors, Release 6,” Technical Report, U.S. Department of Agriculture, Beltsville, MD 2007.
- , “FoodData Central,” 2019.
- Vincent, A., F. Grande, E. Compaoré, G. Amponsah Annor, P. A. Addy, L. C. Aburime, D. Ahmed, A. M. Bih Loh, S. Dahdouh Cabia, N. Deflache, F. M. Dembélé, B. Dieudonné, O. B. Edwige, H. N. Ene-Obong, N. Fanou Fogny, M. Ferreira, J. Omaghomi Jemide, P. C. Kouebou, C. Muller, S. Nájera Espinosa,**

**F. Ouattara, D. Rittenschober, H. Schönfeldt, B. Stadlmayr, M. van Deventer, A. Razikou Yiagnigni, and U. R. Charrondière**, *FAO/INFOODS Food Composition Table for Western Africa (2019) User Guide & Condensed Food Composition Table*, Rome, Italy: Food and Agriculture Organization of the United Nations (FAO), 2020.

## B. Appendix Figures




Figure B1. **Likelihood of consumption, by food group.** HH: household.  
Back to [Section 3](#).




Figure B2. Average share of total energy intake, by food group.  
Back to [Section 5](#).




Figure B3. Distribution of the intake of energy and nutrients. Daily intake per adult equivalent unit.  
Back to [Section 5](#).




Figure B4. **Average share of total energy intake, by food group and source of consumption.** Back to [Section 5](#).




Figure B5. **Agricultural season calendar.** Based on Famine Early Warning Systems Network (FEWSNET). The growing season includes both planting and growing. Back to [Section 3](#).




Figure B6. **Percentage of planted plots by month.** Based on 39,030 plots from sample panel farm HH. Representing all months between the reported planting start date and the reported harvest start date for each plot.  
 Back to [Section 3](#).




Figure B7. **Temperature distribution of growing season days, by region.** Based on the sample farm panel households, i.e., 9,006 observations, 2,832 HH, within 392 enumeration areas (clusters), over 2010-2023.  
 Back to [Section 3](#).




Figure B8. **Optimal GDD/HDD threshold using an iterative regression approach and effect on total yield.** Total yield is measured in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional growing season HDD. Conley spatial HAC standard errors (in parentheses).

Back to [Section 5](#).




Figure B9. **Effect of HDD on consumption likelihood, by food group.** Each food group represents a dummy for household consumption in the last 7 days, but coefficients have been divided by the respective sample means, thus coefficients can be interpreted as the relative percentage change in the likelihood of consumption for one additional HDD. Conley spatial HAC standard errors are used to build the 95% confidence intervals.

Back to [Section 5](#).




Figure B10. **Effect of HDD on energy intake, by food group.** Each food group is expressed as the logarithm of total energy intake, thus coefficients can be interpreted as the relative percentage change for one additional HDD. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors are used to build the 95% confidence intervals.

Back to [Section 5](#).




Figure B11. **Temperatures and livestock birth, purchase, sale, and death.** Poisson pseudo-maximum likelihood estimations using the computationally efficient estimator for Poisson regressions using high-dimensional fixed effects developed by [Correia \(2016\)](#). Coefficients are exponentiated and presented as incidence rate ratios (IRR). Reference bin < 31C. Weather over the last 12 months (number of days in each bin). Mean number of annual days with average temperature > 35C: 0.6. 95% confidence intervals are built using robust clustered standard errors at the households level.

Back to [Section 5](#).




Figure B12. **Effect of HDD on household nutrient adequacy, by survey months.** Each nutrient adequacy represents a dummy equal to one if a household meets the intake requirement at various thresholds (60%, 80%, and 100%). Coefficients have been divided by the respective dependent variable sample mean, and can be interpreted as the relative percentage change in the likelihood of meeting the requirement for one additional HDD. Conley spatial HAC standard errors are used to build the 95% confidence intervals.

Back to [Section 5](#).




Figure B13. **Falsification tests.** Adequacy indicators are set to 80% adequacy levels. Distribution of the estimated coefficients resulting from the falsification tests using random weather allocation. The vertical dotted line in each subplot reflects the respective coefficient estimate of HDD for the main specification. Adq: adequacy. HDDS: Household dietary diversity score. prot: protein.

Back to [Section 5](#).




Figure B14. **Robustness: Effect of HDD on energy intake, by food group.** Total energy intake is expressed in kilocalories and the coefficient is divided by the mean energy intake, thus coefficients in this figure can be interpreted as the relative percentage change for one additional HDD. This provides a robustness to Figure B10, where the outcome variable is expressed as logarithms. Conley spatial HAC standard errors are used to build the 95% confidence intervals. Back to Section 5.

## C. Appendix Tables

	2010	2012	2015	2018	2023	All
Panel farm HH ( $\geq 3$ waves)	2,609	2,599	2,628	968	935	9,739
Strictly positive ag output & area	2,472	2,517	2,530	928	876	9,323
Strictly positive total energy intake	2,398	2,343	2,489	916	860	9,006
<b>Final sample</b>	<b>2,398</b>	<b>2,343</b>	<b>2,489</b>	<b>916</b>	<b>860</b>	<b>9,006</b>
% of panel sample	(91.9%)	(90.2%)	(94.7%)	(94.6%)	(92.0%)	(92.5%)

Table C1. **Sample selection.** A partial refresh of the panel sample was undertaken in wave 4. The representativeness of the survey was impacted during wave 4 due to security issues, which prevented data collection in some locations. The survey does not track “split-off” households. HH: household.

Back to [Section 3](#).

	January	February	March	April	Total
Wave 1	0	41	2,296	61	2,398
Wave 2	0	373	1,964	6	2,343
Wave 3	0	597	1,879	13	2,489
Wave 4	902	14	0	0	916
Wave 5	0	797	63	0	860
Total	902	1,822	6,202	80	9,006

Table C2. **Distribution of interview months, by survey wave.**

Back to [Section 5](#).

	Total yield (kg/ha)	Total harvest (kg)	Total area (ha)
GDD in growing season	-0.0018*** (0.0004)	-0.0017*** (0.0005)	0.0000 (0.0004)
HDD in growing season	-0.0174*** (0.0041)	-0.0154*** (0.0047)	0.0020 (0.0030)
Precipitations	0.0077** (0.0035)	0.0055** (0.0028)	-0.0022 (0.0024)
Precipitations <sup>2</sup> ( $\times 10^{-3}$ )	-0.0157** (0.0077)	-0.0137** (0.0059)	0.0021 (0.0055)
Household FE	Yes	Yes	Yes
Year FE	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes
R <sup>2</sup>	0.506	0.602	0.727
Observations	9,006	9,006	9,006

Table C3. **Growing season weather, agricultural productivity, output, and land area.** Total yield, harvest, and land area are measured in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional growing season HDD. Conley spatial HAC standard errors (in parentheses). FE: fixed effects.

Back to [Section 5](#).

	Energy	Fat	Carbs	Protein
GDD in growing season	0.0001 (0.0002)	0.0000 (0.0002)	0.0001 (0.0003)	0.0001 (0.0002)
HDD in growing season	-0.0002 (0.0015)	0.0002 (0.0012)	-0.0004 (0.0018)	-0.0005 (0.0015)
Household FE	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.554	0.517	0.530	0.605
Observations	9,006	9,006	9,006	9,006

Table C4. **Temperatures and energy and macronutrient intake.** Expressed as logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Conley spatial HAC standard errors (in parentheses). FE: fixed effects.

Back to [Section 5](#).

	Total	Purchased	Own-prod	Gifted
GDD in growing season	0.0001 (0.0001)	-0.0001 (0.0001)	0.0002 (0.0002)	-0.0001 (0.0002)
HDD in growing season	-0.0002 (0.0008)	-0.0015 (0.0013)	0.0002 (0.0016)	-0.0020 (0.0015)
Household FE	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.554	0.558	0.905	0.895
Observations	9,006	9,006	9,006	9,006

Table C5. **Temperatures and energy intake, by source of consumption.** Expressed as logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors (in parentheses). FE: fixed effects. prod: production.

Back to [Section 5](#).

	Education	Housing	Comm & transport	Clothing	Recreation
GDD in growing season	-0.0003* (0.0002)	-0.0008*** (0.0002)	-0.0005** (0.0002)	-0.0006** (0.0003)	-0.0001 (0.0003)
HDD in growing season	-0.0013 (0.0017)	-0.0094*** (0.0017)	-0.0065*** (0.0017)	-0.0063*** (0.0021)	0.0004 (0.0025)
Household FE	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.808	0.728	0.863	0.570	0.662
Observations	8,146	8,146	8,146	8,146	8,146

Table C6. **Temperatures and non-food expenditure.** Housing includes utilities and household services. Comm: communication. Clothing includes housing goods. Recreation includes other miscellaneous expenses. Expenditure is expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Annualized expenditure in USD 2020 (originally, food: last 7 days; non-food: last month or last 12 months). Expenditure information is missing for wave 5. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors (in parentheses). FE: fixed effects. HDDS: Household dietary diversity score.

Back to [Section 5](#).

	Total	Purchased	Own-prod	Gifted
GDD	0.0004* (0.0002)	0.0001 (0.0003)	0.0005 (0.0003)	0.0000 (0.0002)
GDD × Mar-Apr	-0.0004*** (0.0001)	-0.0003*** (0.0001)	-0.0005*** (0.0001)	-0.0002** (0.0001)
HDD	0.0015 (0.0015)	-0.0007 (0.0022)	0.0026 (0.0026)	-0.0015 (0.0019)
HDD × Mar-Apr	-0.0015** (0.0007)	0.0002 (0.0010)	-0.0034** (0.0013)	-0.0009 (0.0008)
Household FE	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.563	0.561	0.905	0.895
Observations	9,006	9,006	9,006	9,006

Table C7. **Temperatures and energy intake, by source of consumption and survey months.** Variables are expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors (in parentheses). FE: fixed effects. Back to [Section 5](#).

	Cereals	Roots	Veg	Fruits	Pulses
GDD	0.0004** (0.0002)	0.0001 (0.0003)	0.0001 (0.0002)	-0.0001 (0.0001)	0.0001 (0.0001)
GDD × Mar-Apr	-0.0003*** (0.0001)	0.0000 (0.0001)	-0.0003*** (0.0001)	0.0000 (0.0000)	-0.0002*** (0.0001)
HDD	0.0033* (0.0018)	-0.0009 (0.0018)	-0.0011 (0.0013)	-0.0019** (0.0007)	0.0011 (0.0010)
HDD × Mar-Apr	-0.0021** (0.0009)	0.0000 (0.0006)	-0.0012* (0.0006)	0.0000 (0.0002)	-0.0010* (0.0005)
Household FE	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.982	0.974	0.851	0.961	0.984
Observations	8,090	8,090	8,090	8,090	8,090

Table C8. **Temperatures and energy intake from main own-produced foods, by survey months.** Variables are expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors (in parentheses). FE: fixed effects. Back to [Section 5](#).

	HDDS	Adq prot	Adq iron	Food exp	Non-food exp	Share sold
GDD	-0.0004 (0.0004)	0.0000 (0.0002)	0.0000 (0.0002)	0.0000 (0.0003)	-0.0006*** (0.0001)	-0.0001 (0.0001)
GDD × Mar-Apr	-0.0001 (0.0002)	-0.0001*** (0.0000)	-0.0002*** (0.0001)	-0.0002* (0.0001)	-0.0000 (0.0001)	-0.0000 (0.0000)
HDD	-0.0028 (0.0031)	-0.0014 (0.0010)	-0.0016* (0.0010)	-0.0030* (0.0018)	-0.0088*** (0.0013)	-0.0014** (0.0006)
HDD × Mar-Apr	-0.0030** (0.0015)	-0.0010** (0.0004)	-0.0009* (0.0005)	-0.0007 (0.0010)	0.0017** (0.0008)	0.0001 (0.0004)
Household FE	Yes	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.621	0.518	0.522	0.627	0.766	0.512
Observations	8,090	8,090	8,090	7,230	7,230	5,330
Mean Y	5.521	0.826	0.752	.	.	0.197

Table C9. **Robustness: Temperatures and dietary diversity, nutrient adequacy, expenditure, and commercialization, by survey months.** This table is a robustness analysis for Table 4, excluding wave 4. Adequacy indicators are set to 80% adequacy levels. Coefficients on HDDS can be interpreted as the absolute change in the HDDS for one additional HDD. Coefficients on adequacy indicators can be interpreted as percentage-point change in the likelihood for one additional HDD. Expenditure is expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Annualized expenditure in USD 2020 (originally, food: last 7 days; non-food: last month or last 12 months). Food expenditure represents purchased food. Expenditure information is missing for wave 5. I set the dependent variable (logarithm) to zero for all zero consumption values and include a dummy variable in the regression to account for this data transformation. Information on sold harvested quantities is missing for 33.7% of households (either missing or set to missing because it is higher than harvested quantities or negative). Conley spatial HAC standard errors (in parentheses). Adq: Adequacy. FE: fixed effects. HDDS: Household dietary diversity score. Back to Section 5.

	HDDS	Adq prot	Adq iron	Food exp	Non-food exp	Share sold
HDD	-0.0053* (0.0031)	-0.0020** (0.0008)	-0.0025*** (0.0009)	-0.0036** (0.0017)	-0.0069*** (0.0015)	-0.0011* (0.0006)
HDD × HH livestock	-0.0003 (0.0015)	-0.0003 (0.0003)	-0.0001 (0.0004)	-0.0003 (0.0007)	-0.0009 (0.0006)	-0.0008*** (0.0003)
HDD	-0.0065* (0.0035)	-0.0016 (0.0010)	-0.0014 (0.0011)	-0.0041** (0.0019)	-0.0077*** (0.0016)	-0.0018** (0.0007)
HDD × ln(dist pop center)	0.0004 (0.0005)	-0.0001 (0.0001)	-0.0002* (0.0001)	0.0001 (0.0002)	0.0000 (0.0002)	0.0001 (0.0001)
HDD	-0.0064** (0.0030)	-0.0021** (0.0008)	-0.0023** (0.0010)	-0.0036** (0.0017)	-0.0075*** (0.0014)	-0.0017*** (0.0006)
HDD × HH non-farm enterprise	0.0019 (0.0014)	-0.0003 (0.0003)	-0.0003 (0.0004)	-0.0003 (0.0008)	-0.0001 (0.0006)	-0.0001 (0.0003)
HDD	-0.0051* (0.0028)	-0.0022*** (0.0008)	-0.0026*** (0.0009)	-0.0040** (0.0016)	-0.0076*** (0.0014)	-0.0020*** (0.0006)
HDD × Head wage job (last 7 days)	0.0001 (0.0019)	0.0002 (0.0004)	0.0011** (0.0005)	-0.0006 (0.0010)	0.0007 (0.0011)	0.0012*** (0.0004)
Mean Y	5.576	0.829	0.750	.	.	0.199

Table C10. **Effect of HDD on dietary diversity, nutrient adequacy, expenditure, and commercialization, by potential moderators.** This table displays the results from 24 regressions, under the main specification. Adequacy indicators are set to 80% adequacy levels. Coefficients on HDDS can be interpreted as the absolute change in the HDDS for one additional HDD. Coefficients on adequacy indicators can be interpreted as percentage-point change in the likelihood for one additional HDD. Expenditure is expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Annualized expenditure in USD 2020 (originally, food: last 7 days; non-food: last month or last 12 months). Food expenditure represents purchased food. Expenditure information is missing for wave 5. Information on sold harvested quantities is missing for 33.7% of households (either missing or set to missing because it is higher than harvested quantities or negative). HH livestock, HH non-farm enterprise, and head wage job are dummy variables equal to one if the household manages at least one livestock head, ran at least one non-farm enterprise in last 12 months, was employed in a wage job in the last 7 days. The distance to the nearest population center with more than 20,000 inhabitants is in log kilometers. Conley spatial HAC standard errors (in parentheses). Adq: Adequacy. dist: distance. edu: education. HDDS: Household dietary diversity score. H: household. pop: population. prot: protein.

Back to [Section 5](#).

	Harvest (kg)	HDDS	Adq prot	Adq iron
Distance 50km, time lag 1 year	(0.0038)***	(0.0025)**	(0.0007)***	(0.0008)***
Distance 50km, time lag 5 years	(0.0038)***	(0.0026)**	(0.0007)***	(0.0008)***
Distance 50km, time lag 10 years	(0.0039)***	(0.0027)*	(0.0007)***	(0.0009)***
Distance 100km, time lag 1 year	(0.0046)***	(0.0028)*	(0.0008)***	(0.0009)***
Distance 100km, time lag 5 years	(0.0047)***	(0.0028)*	(0.0008)***	(0.0009)***
Distance 100km, time lag 10 years	(0.0047)***	(0.0029)*	(0.0008)***	(0.0009)***
Distance 200km, time lag 1 year	(0.0056)***	(0.0030)*	(0.0008)***	(0.0009)***
Distance 200km, time lag 5 years	(0.0056)***	(0.0031)*	(0.0009)***	(0.0009)***
Distance 200km, time lag 10 years	(0.0057)***	(0.0032)*	(0.0009)**	(0.0009)***

Table C11. **Spatial-HAC standard errors sensitivity.** Adequacy indicators are set to 80% adequacy levels. Conley spatial HAC standard errors on the HDD coefficient for the main specification. Adq: Adequacy. HDDS: Household dietary diversity score. prot: protein. Back to [Section 5](#).

	Maize	Potato	Onion	Banana	Beef	Egg	Fish	Horsebean	Milk
GDD in growing season	0.0001 (0.0003)	-0.0004 (0.0020)	-0.0008 (0.0012)	0.0003 (0.0008)	-0.0006 (0.0012)	0.0027*** (0.0008)	0.0008 (0.0007)	-0.0004 (0.0006)	-0.0018 (0.0023)
HDD in growing season	-0.0021 (0.0035)	0.0106 (0.0222)	-0.0102 (0.0096)	-0.0156* (0.0093)	-0.0061 (0.0118)	0.0296** (0.0122)	-0.0075 (0.0065)	-0.0050 (0.0050)	-0.0147 (0.0167)
Cluster FE	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.771	0.633	0.408	0.606	0.519	0.721	0.553	0.717	0.500
Clusters	110	32	131	120	328	88	252	89	37
Observations	283	73	382	333	891	219	693	231	89

Table C12. **Temperatures and food prices.** Variables are expressed in logarithms, thus coefficients can be interpreted as the relative percentage change for one additional HDD. Conley spatial HAC standard errors (in parentheses).

Back to [Section 5](#).

	< 5 yo	15-65 yo	> 65 yo
GDD in growing season	-0.0001 (0.0001)	0.0001 (0.0001)	-0.0000 (0.0000)
HDD in growing season	-0.0005 (0.0006)	0.0010 (0.0007)	-0.0001 (0.0001)
Household FE	Yes	Yes	Yes
Year FE	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes
R <sup>2</sup>	0.622	0.792	0.578
Observations	8,949	8,941	8,949

Table C13. **Temperatures and household composition.** Each age group is expressed as the logarithm of total household member in that age range, thus coefficients can be interpreted as the relative percentage change for one additional HDD. I set the dependent variable (logarithm) to zero for all zero values and include a dummy variable in the regression to account for this data transformation. Conley spatial HAC standard errors (in parentheses).

Back to [Section 5](#).

	Share of total area (ext. margin)				Intensive margin (kg)
	Mixed cropping	Pesticides	Organic fert.	Inorganic fert.	Inorganic fert.
GDD in growing season	0.0001 (0.0001)	-0.0001 (0.0001)	-0.0003*** (0.0001)	0.0002 (0.0002)	-0.0009*** (0.0002)
HDD in growing season	0.0006 (0.0007)	0.0004 (0.0008)	-0.0000 (0.0010)	0.0008 (0.0011)	-0.0046*** (0.0017)
Household FE	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.550	0.529	0.509	0.598	0.853
Observations	9,006	9,006	9,006	9,006	9,006
Mean Y	0.678	0.180	0.116	0.380	.

Table C14. **Temperatures and farm household productive response.** Conley spatial HAC standard errors (in parentheses). Ext.: extensive margin. FE: fixed effects. Int.: intensive margin.

Back to [Section 5](#).

	Total	Family		Hired	
	Int.	Ext.	Int.	Ext.	Int.
GDD in growing season	0.0009 (0.0009)	-0.0001** (0.0000)	0.0010 (0.0009)	0.0001 (0.0002)	0.0005* (0.0003)
HDD in growing season	0.0048 (0.0057)	-0.0004 (0.0002)	0.0059 (0.0059)	-0.0006 (0.0013)	0.0016 (0.0020)
Household FE	Yes	Yes	Yes	Yes	Yes
Year FE	Yes	Yes	Yes	Yes	Yes
Month-of-int FE	Yes	Yes	Yes	Yes	Yes
Precipitations	Yes	Yes	Yes	Yes	Yes
R <sup>2</sup>	0.690	0.391	0.704	0.537	0.844
Observations	9,006	9,006	9,006	9,006	9,006
Mean Y	.	0.985	.	0.679	.

Table C15. **Temperatures and farm labor during the agricultural season.** Conley spatial HAC standard errors (in parentheses). Ext.: extensive margin. FE: fixed effects. Int.: intensive margin. SOB: small-own business.

Back to [Section 5](#).